

Fact Sheet

PSYCHOSIS

What is the Issue?

Psychosis is a condition which impairs a person's sense of reality. This may be from a mental illness such as schizophrenia or bipolar disorder (manic depression). At times, symptoms of psychosis can also be caused by brain injury or infection, or alcohol or drug misuse.

Most young people recover from psychosis to live full and satisfying lives.

No-one knows for sure what causes psychotic illnesses such as schizophrenia or bipolar disorder. The first signs usually appear in late adolescence and early adulthood and may begin gradually.

The earliest phase of a psychotic disorder can last from a few weeks, to up to two years. Initially, vague changes may occur in the way a person thinks, feels and behaves. At first, these signs may be barely noticeable. A young person in this phase may experience confused thinking, withdrawal from previously enjoyed activities and deterioration in school performance.

Psychosis may affect a young person's ability to concentrate, or their thoughts might seem faster or slower than usual. A young person experiencing these changes may feel very confused and distressed by what is happening.

Psychosis can happen to anyone, but like other illnesses, it can be treated.

Early warning signs

The early signs of psychosis may be difficult to detect and can be mistaken for normal changes of adolescence. Noticeable and persistent changes in the young person's behaviour and moods should not be ignored.

The early warning signs in young people are:

- *Changes in thinking* – difficulty in concentrating, poor memory, preoccupation with odd ideas, increased suspiciousness.
- *Changes in mood* – lack of emotional response, rapid mood changes, inappropriate moods.
- *Changes in behaviour* – odd or unusual behaviour.
- *Physical changes* – sleep disturbance or excessive sleep and loss of energy.
- *Social changes* – withdrawal or isolation from friends and family.
- *Changes in functioning* – decline in school or work performance.

The term psychosis refers to a group of disorders which impair a person's sense of reality and may lead to changes in their mood. A person's perceptions can be affected by hallucinations. As well, their thinking may seem confused, slowed down or speeded up. They may have persistent false beliefs known as delusions. The symptoms vary for each person and may change over time. Psychosis may begin in adolescence. It can happen to anyone but like most other illnesses, it can be treated.

Helping young people with psychosis

Young people with first onset psychosis may be reluctant to seek treatment because they don't think anything is wrong or they hope their symptoms will go away. Help is needed for the young person to find out what is happening and what treatment is required. Families also need support to understand what is happening and how they can help.

It is important to get help as early as possible as this improves recovery for young people. The impact of psychosis can be very traumatic for the young person and can disrupt their life at a critical developmental stage. The longer it takes to receive effective treatment, the longer it is likely to take for symptoms to go away. Delayed help also increases the risk of further episodes.

Source: Child and Youth Mental Health Service: Adapted from NSW Health Brochures
Last Updated: September 2002
Review Date: September 2005

Fact Sheet

Mental health professionals treat most young people with psychosis in the community. This minimises distress and disruption for the young person and their family. Having someone to talk to is an important part of treatment. Mental health workers understand what is happening and can provide reassurance and information for the young person and their family.

It is a good idea to arrange a check-up with your doctor to ensure that there are no other serious physical or emotional problems which may require treatment.

Medication is often a part of the treatment of psychosis to assist recovery and prevent further episodes. A psychiatrist can determine if medication is needed. There are several different types of medication that may be recommended and young people usually start on very low doses.

Does anyone recover from psychosis?

Recovery varies for each person, with some returning quickly to their usual lifestyle, while others may take months or longer to get back on track. Most young people will recover from psychosis to live full and satisfying lives.

Family and friends may help the young person's recovery by:

- Finding out about the signs and effects of psychosis.
- Offering support and encouragement.
- Allowing the young person time for recovery and return to school, university, work or job training.
- Obtaining support for themselves.

The first point of contact can be:

- Your child's general practitioner, teacher, guidance officer, school counsellor or school health nurse.
- If more specialised assessment or intervention is required, general practitioners, schools or other health professionals can make a referral to a Child and Youth Mental Health Service (CYMHS) – for local clinics, look under Health in the White Pages telephone directory.
- Your general practitioner may refer you to other specialists who work with children and adolescents such as a private psychiatrist or psychologist.
- In an emergency contact your local hospital Emergency Department.

Other support organisations:

Kids Help Line

Phone: 1800 551 800

24 hours, every day of the year

ARAFMI - Association of Relatives & Friends of the Mentally Ill

PO Box 248

NEW FARM QLD 4005

Phone: (07) 3254 1881 (24 hours)

Lifeline

Phone: 13 11 14

24 hours, every day of the year

Schizophrenia Fellowship of South Queensland

95 Arthur Street

FORTITUDE VALLEY QLD 4006

Phone: (07) 3358 4424 (office hours)

Source: Child and Youth Mental Health Service: Adapted from NSW Health Brochures

Last Updated: September 2002

Review Date: September 2005