Queensland Government	
--------------------------	--

Fractured Hip/	G
Hemi-Arthroplasy	A
Facility:	D
A. Interpreter / cultural needs	
An Interpreter Service is required?	Yes No
If Yes, is a qualified Interpreter present?	Yes No

B. Condition and treatment

A Cultural Support Person is required?

The doctor has explained that you have the following condition: (Doctor to document in patient's own words)

If Yes, is a Cultural Support Person present? Yes No

This condition requires the following procedure.	
Doctor to document - include site and/or side whe	re
relevant to the procedure)	

Left hip	☐ Yes ☐ No
Right hip	Yes No

The following will be performed:

A fractured hip hemi-arthroplasty procedure is the replacement of half of the fractured hip joint with an artificial hip joint - a hemi-arthroplasty or Thompson's prosthesis.

C. Risks of a fractured hip/ hemiarthroplasy

There are risks and complications with this procedure. They include but are not limited to the following.

General risks:

- Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

Given name(s):			
Address:			
Date of birth:	Sex: M	F	

(Affix identification label here)

Specific risks:

URN:

Yes No

Family name: Given nam Address:

- Blood clots can form in the legs. Drugs and compression stockings are usually used to help prevent this. The clots can break off and travel to the lungs and can cause death.
- Infection after hip replacement. This may require further surgery and possibly the new hip to be removed.
- A tube into the bladder at the time of surgery. In which case, the bladder may not empty after the tube is removed after surgery.
- The bowel may be paralysed or blocked after surgery. This may cause a build up of fluid in the bowel with bloating and vomiting. This may need further treatment.
- The hip joint can dislocate. The hip can usually be put back in without surgery. A brace may be worn for some weeks.
- The joint may break during surgery. A second operation will be required to repair the break.
- The nerves around the hip joint may be injured during the surgery. This causes pain, and/ or paralysis, which can cause permanent disability.
- The artificial joint may loosen which can happen over time.
- The leg length may be different to the unoperated side. Further treatment will be required to correct the length.

ROCE

CONSENT


- Possible bleeding into the wound after the surgery. This may be painful and require surgical drainage, or become infected, needing antibiotics.
- The wound may not heal properly and can become red, thickened and painful. This can be disfiguring.
- There is a risk that the leg may have to be amputated due to poor blood supply or infection.
- Infection can spread to the artificial hip joint from other areas in the body. To prevent this, you will need antibiotics before other procedures and dental work.
- The hip joint may fail within five years of surgery. Further surgery will be required to correct the cause of the failure.
- -This consent document continues on page 2 -



	(Affix identification labe	l here)		
URN:				
Family name:				
Given name(s):				
Address:				
Date of birth:		Sex: M	F	

Fractured Hip/ Hemi-Arthroplasy	G
Facility:	D
D. Significant risks and procedure options	
Doctor to document in space provided. Continue in Medical Record if necessary.)	
E. Risks of not having this procedure	
Doctor to document in space provided. Continue in Medical Record if necessary.)	
,,,	
F. Anaesthetic	
This procedure may require an anaesthetic. (Doctor	to
document type of anaesthetic discussed)	

This consent document continues on page 3





Fractured Hip/ **Hemi-Arthroplasy**

Facility:

G. Patient consent

I acknowledge that the doctor has explained;

- my medical condition and the proposed procedure, including additional treatment if the doctor finds something unexpected. I understand the risks, including the risks that are specific to
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- other relevant procedure/treatment options and their associated risks.
- my prognosis and the risks of not having the procedure.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- the procedure may include a blood transfusion.
- tissues and blood may be removed and could be used for diagnosis or management of my condition, stored and disposed of sensitively by the hospital.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor or my Acute Resuscitation Plan.
- a doctor other than the Consultant may conduct the procedure. I understand this could be a doctor undergoing further training.

I have been given the following Patient Information Sheet/s:

- About Your Anaesthetic OR **Epidural & Spinal Anaesthesia Fractured Hip Hemi-Arthroplasty Blood & Blood Products Transfusion**
- I was able to ask questions and raise concerns with the doctor about my condition, the proposed procedure and its risks, and my treatment options. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time, including after I have signed this form but, preferably following a discussion with my doctor.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.

On the basis of the above statements,

(Affix identification label here)			
URN:			
Family name:			
Given name(s):			
Address:			
Date of birth:		Sex: M	F I

I reques	t to have the procedure
Name of Pa	atient:
Signature:	
Date:	
Consent	s who lack capacity to provide consent must be obtained from a substitute decision in the order below.
Does the (AHD)?	patient have an Advance Health Directive
☐ Yes ►	Location of the original or certified copy of the AHD:
□ No ►	Name of Substitute Decision Maker/s: Signature:
	Relationship to patient:
	Date: PH No:
	Source of decision making authority (tick one): Tribunal-appointed Guardian
	Attorney/s for health matters under Enduring Power of Attorney or AHD
	Statutory Health Attorney If none of these, the Adult Guardian has provided
	consent. Ph 1300 QLD OAG (753 624)

H. Doctor/delegate Statement

I have explained to the patient all the above points under the Patient Consent section (G) and I am of the opinion that the patient/substitute decisionmaker has understood the information.

Name of Doctor/delegate: Designation: ____ Signature:.. Date:

Interpreter's statement

I have given a sight translation in

(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

Signature:

ame of	
iterpreter:	
	•

Date:



Consent Information - Patient Copy Fractured Hip/ Hemi-Arthroplasy

What do I need to know about this procedure??

A fractured hip hemi-arthroplasty procedure is the replacement of half of the fractured hip joint with an artificial hip joint - a hemi-arthroplasty or Thompson's prosthesis.

2. My anaesthetic

This procedure will require an anaesthetic.

See About Your Anaesthetic OR Epidural and Spinal Anaesthetic for information about the anaesthetic and the risks involved. If you have any concerns, discuss these with your doctor.

If you have not been given an information sheet, please ask for one.

3. What are the risks of this specific procedure?

There are risks and complications with this procedure. They include but are not limited to the following.

- · Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- · Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

Specific risks:

- · Blood clots can form in the legs. Drugs and compression stockings are usually used to help prevent this. The clots can break off and travel to the lungs and can cause death.
- Infection after hip replacement. This may require further surgery and possibly the new hip to be removed.
- A tube into the bladder at the time of surgery. In which case, the bladder may not empty after the tube is removed after surgery.
- The bowel may be paralysed or blocked after surgery. This may cause a build up of fluid in the bowel with bloating and vomiting. This may need further treatment.
- The hip joint can dislocate. The hip can usually be put back in without surgery. A brace may be worn for some weeks.

- The joint may break during surgery. A second operation will be required to repair the break.
- The nerves around the hip joint may be injured during the surgery. This causes pain, and/ or paralysis, which can cause permanent disability.
- The artificial joint may loosen which can happen over time.
- The leg length may be different to the unoperated side. Further treatment will be required to correct the length.
- · Possible bleeding into the wound after the surgery. This may be painful and require surgical drainage, or become infected, needing antibiotics.
- The wound may not heal properly and can become red, thickened and painful. This can be disfiguring.
- There is a risk that the leg may have to be amputated due to poor blood supply or infection.
- Infection can spread to the artificial hip joint from other areas in the body. To prevent this, you will need antibiotics before other procedures and dental work.
- The hip joint may fail within five years of surgery. Further surgery will be required to correct the cause of the failure.

Notes to talk to my doctor about:						