

TABLE 6.01

**MOTHERS DELIVERING IN QUEENSLAND, 1998**  
**ONSET OF LABOUR BY PLURALITY OF PREGNANCY**

Onset of labour	Plurality of pregnancy					
	Singleton		Multiple		Total	
	No.	%	No.	%	No.	%
Spontaneous - not augmented	16,998	36.3	179	26.1	17,177	36.2
Spontaneous - augmented	12,575	26.9	93	13.5	12,668	26.7
Induced	11,323	24.2	161	23.4	11,484	24.2
No labour	5,864	12.5	254	37.0	6,118	12.9
Not stated	3	0.0	-	-	3	0.0
<b>Total</b>	<b>46,763</b>	<b>100.0</b>	<b>687</b>	<b>100.0</b>	<b>47,450</b>	<b>100.0</b>

**TABLE 6.02****MOTHERS DELIVERING IN QUEENSLAND, 1998  
ONSET OF LABOUR BY AGE OF MOTHER**

Onset of labour	Age of mother							
	Under 20		20 - 34		35 and over		Total	
	No.	%	No.	%	No.	%	No.	%
Spontaneous - not augmented	1,226	39.7	13,727	36.3	2,224	34.1	17,177	36.2
Spontaneous - augmented	1,030	33.4	10,266	27.1	1,372	21.1	12,668	26.7
Induced	715	23.2	9,238	24.4	1,531	23.5	11,484	24.2
No labour	114	3.7	4,615	12.2	1,389	21.3	6,118	12.9
Not stated	-	-	3	0.0	-	-	3	0.0
<b>Total</b>	<b>3,085</b>	<b>100.0</b>	<b>37,849</b>	<b>100.0</b>	<b>6,516</b>	<b>100.0</b>	<b>47,450</b>	<b>100.0</b>

**TABLE 6.03      MOTHERS WITH INDUCED LABOURS, QUEENSLAND, 1998**  
**SELECTED REASON FOR INDUCTION**

Reason for induction	Mothers with induced labour	
	No.	%
Prolonged pregnancy	4,175	36.4
Hypertension complicating pregnancy	2,008	17.5
Unspecified psychosocial circumstances	1,356	11.8
Premature/prolonged rupture of membranes	1,245	10.8
Poor fetal growth	464	4.0
Diabetes complicating pregnancy	293	2.6
Previous poor obstetric history	201	1.8
Spurious/false labour	175	1.5
Other fetal/placental problem	136	1.2
Intrauterine fetal death	115	1.0
Other reasons	1,316	11.5
<b>Total</b>	<b>11,484</b>	<b>100.0</b>

**TABLE 6.04 BIRTHS, QUEENSLAND, 1998  
PRESENTATION BY PLURALITY OF PREGNANCY**

Presentation	Plurality of pregnancy					
	Singleton		Multiple		Total	
	No.	%	No.	%	No.	%
Vertex	44,459	95.1	914	65.3	45,373	94.2
Face	50	0.1	2	0.1	52	0.1
Brow	46	0.1	3	0.2	49	0.1
Breech <sup>(a)</sup>	1,965	4.2	419	29.9	2,384	4.9
Other	239	0.5	62	4.4	301	0.6
Not stated	4	0.0	-	-	4	0.0
<b>Total</b>	<b>46,763</b>	<b>100.0</b>	<b>1,400</b>	<b>100.0</b>	<b>48,163</b>	<b>100.0</b>

(a) Breech deliveries in which forceps were applied to the after-coming head were included in the breech' classification and not classified as 'forceps' deliveries.

**TABLE 6.05 BIRTHS, QUEENSLAND, 1998  
METHOD OF DELIVERY BY PLURALITY OF PREGNANCY**

Method of delivery	Plurality of pregnancy					
	Singleton		Multiple		Total	
	No.	%	No.	%	No.	%
Spontaneous vertex	31,902	68.2	378	27.0	32,280	67.0
Forceps	2,103	4.5	77	5.5	2,180	4.5
Vacuum extraction	2,043	4.4	55	3.9	2,098	4.4
LSCS	10,331	22.1	769	54.9	11,100	23.0
Classical CS	82	0.2	6	0.4	88	0.2
Breech	259	0.6	109	7.8	368	0.8
Other	41	0.1	6	0.4	47	0.1
Not stated	2	0.0	-	-	2	0.0
<b>Total</b>	<b>46,763</b>	<b>100.0</b>	<b>1,400</b>	<b>100.0</b>	<b>48,163</b>	<b>100.0</b>

**TABLE 6.06 BIRTHS, QUEENSLAND, 1998  
METHOD OF DELIVERY BY AGE OF MOTHER**

Method of delivery	Age of mother (years)							
	Under 20		20 - 34		35 and over		Total	
	No.	%	No.	%	No.	%	No.	%
Spontaneous vertex	2,467	79.4	25,900	67.4	3,913	58.8	32,280	67.0
Forceps	123	4.0	1,772	4.6	285	4.3	2,180	4.5
Vacuum extraction	106	3.4	1,748	4.6	244	3.7	2,098	4.4
LSCS	383	12.3	8,576	22.3	2,141	32.2	11,100	23.0
Classical CS	4	0.1	63	0.2	21	0.3	88	0.2
Breech	21	0.7	300	0.8	47	0.7	368	0.8
Other	2	0.1	39	0.1	6	0.1	47	0.1
Not stated	-	-	2	0.0	-	-	2	0.0
<b>Total</b>	<b>3,106</b>	<b>100.0</b>	<b>38,400</b>	<b>100.0</b>	<b>6,657</b>	<b>100.0</b>	<b>48,163</b>	<b>100.0</b>

TABLE 6.07

## BIRTHS, QUEENSLAND, 1998 - METHOD OF DELIVERY BY PRESENTATION

Method of delivery	Presentation											
	Vertex		Face		Brow		Breech		Other		Total <sup>(a)</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Singleton</b>												
Spontaneous vertex	31,860	71.7	2	4.0	5	10.9	1	0.1	33	13.8	31,902	68.2
Forceps	2,067	4.6	1	2.0	3	6.5	20	1.0	12	5.0	2,103	4.5
Vacuum extraction	2,034	4.6	2	4.0	3	6.5	-	-	4	1.7	2,043	4.4
LSCS	8,445	19.0	26	52.0	31	67.4	1,658	84.4	171	71.5	10,331	22.1
Classical CS	41	0.1	1	2.0	1	2.2	26	1.3	13	5.4	82	0.2
Breech	-	-	-	-	-	-	259	13.2	-	-	259	0.6
Other	12	0.0	18	36.0	3	6.5	1	0.1	6	2.5	41	0.1
Not stated	-	-	-	-	-	-	-	-	-	-	2	0.0
<b>Total</b>	<b>44,459</b>	<b>100.0</b>	<b>50</b>	<b>100.0</b>	<b>46</b>	<b>100.0</b>	<b>1,965</b>	<b>100.0</b>	<b>239</b>	<b>100.0</b>	<b>46,763</b>	<b>100.0</b>
<b>Multiple</b>												
Spontaneous vertex	375	41.0	-	-	-	-	-	-	3	4.8	378	27.0
Forceps	70	7.7	1	50.0	-	-	4	1.0	2	3.2	77	5.5
Vacuum extraction	55	6.0	-	-	-	-	-	-	-	-	55	3.9
LSCS	409	44.7	-	-	3	100.0	307	73.3	50	80.6	769	54.9
Classical CS	1	0.1	-	-	-	-	3	0.7	2	3.2	6	0.4
Breech	-	-	-	-	-	-	105	25.1	4	6.5	109	7.8
Other	4	0.4	1	50.0	-	-	-	-	1	1.6	6	0.4
Not stated	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>914</b>	<b>100.0</b>	<b>2</b>	<b>100.0</b>	<b>3</b>	<b>100.0</b>	<b>419</b>	<b>100.0</b>	<b>62</b>	<b>100.0</b>	<b>1,400</b>	<b>100.0</b>
<b>Total births</b>	<b>45,373</b>		<b>52</b>		<b>49</b>		<b>2,384</b>		<b>301</b>		<b>48,163</b>	

(a) Includes births with a not stated presentation.

**TABLE 6.08 BIRTHS, QUEENSLAND, 1998  
METHOD OF DELIVERY BY METHOD OF INTRAPARTUM ANALGESIA/ANAESTHESIA**

Method of Analgesia/ Anaesthesia <sup>(a)</sup>	Method of delivery									
	Spontaneous		Forceps/ Vacuum		Caesarean		Other		Total <sup>(b)</sup>	
	Vertex				Section					
	No.	%	No.	%	No.	%	No.	%	No.	%
None reported	8,311	25.7	145	3.4	-	-	93	22.4	8,549	17.8
At least one method	23,967	74.2	4,133	96.6	11,188	100.0	320	77.1	39,608	82.2
<i>Nitrous oxide</i>	18,049	75.3	2,497	60.4	1,546	13.8	142	44.4	22,234	56.1
<i>Narcotic(IM/IV)</i>	10,499	43.8	1,803	43.6	1,335	11.9	114	35.6	13,751	34.7
<i>Caudal</i>	49	0.2	53	1.3	37	0.3	7	2.2	146	0.4
<i>Epidural</i>	4,911	20.5	2,675	64.7	5,562	49.7	113	35.3	13,261	33.5
<i>Spinal</i>	331	1.4	213	5.2	5,140	45.9	16	5.0	5,700	14.4
<i>General Anaesthesia</i>	81	0.3	60	1.5	1,408	12.6	14	4.4	1,563	3.9
<i>Other</i>	317	1.3	217	5.3	22	0.2	10	3.1	566	1.4
Not stated	2	0.0	-	-	-	-	2	0.5	6	0.0
<b>Total</b>	<b>32,280</b>	<b>100.0</b>	<b>4,278</b>	<b>100.0</b>	<b>11,188</b>	<b>100.0</b>	<b>415</b>	<b>100.0</b>	<b>48,163</b>	<b>100.0</b>

(a) Reporting of multiple methods of analgesia/anaesthesia permitted.

(b) Includes births with not stated method of delivery.

**TABLE 6.09      MOTHERS DELIVERING BY PRIMARY CAESAREAN SECTION, QUEENSLAND, 1998**  
**LABOUR STATUS BY REASON OF CAESAREAN**


Reason for Caesarean Section	Labour status					
	No labour		Labour		Total	
	No.	%	No.	%	No.	%
Failure to progress	-	-	1,397	36.9	1,397	21.6
Fetal distress	102	3.8	1,086	28.7	1,188	18.3
Breech	817	30.3	343	9.1	1,160	17.9
Hypertensive disorders	325	12.1	66	1.7	391	6.0
Unspecified psychosocial circumstances	275	10.2	28	0.7	303	4.7
Antepartum haemorrhage (including placenta praevia)	212	7.9	79	2.1	291	4.5
Cephalopelvic disproportion	122	4.5	122	3.2	244	3.8
Multiple pregnancy	83	3.1	25	0.7	108	1.7
Other	759	28.2	639	16.9	1,398	21.6
<b>Total</b>	<b>2,695</b>	<b>100.0</b>	<b>3,785</b>	<b>100.0</b>	<b>6,480</b>	<b>100.0</b>

**TABLE 6.10**

**BIRTHS, QUEENSLAND, 1998  
ONSET OF LABOUR BY METHOD OF DELIVERY**

Onset of labour	Method of delivery									
	Spontaneous		Forceps/ Vacuum		Caesarean		Other		Total <sup>(a)</sup>	
	Vertex				Section					
	No.	%	No.	%	No.	%	No.	%	No.	%
Spontaneous - not augmented	14,259	44.2	997	23.3	1,872	16.7	203	48.9	17,331	36.0
Spontaneous - augmented	9,820	30.4	1,678	39.2	1,202	10.7	93	22.4	12,793	26.6
Induced	8,200	25.4	1,603	37.5	1,723	15.4	119	28.7	11,645	24.2
No labour	-	-	-	-	6,391	57.1	-	-	6,391	13.3
Not stated	1	0.0	-	-	-	-	-	-	3	0.0
<b>Total</b>	<b>32,280</b>	<b>100.0</b>	<b>4,278</b>	<b>100.0</b>	<b>11,188</b>	<b>100.0</b>	<b>415</b>	<b>100.0</b>	<b>48,163</b>	<b>100.0</b>

(a) Includes births with not stated method of delivery.




Of the 48,163 babies born in 1998, 41,769 (86.7 per cent) were born after a period of labour.

There were 6,391 babies born via Caesarean section prior to the onset of labour. Of these babies, 2,902 (45.4 per cent) had "Repeat Elective" reported as principal reason for Caesarean section.

**TABLE 6.11 TERM <sup>(a)</sup>, SINGLETON BIRTHS WITH A VERTEX PRESENTATION TO PRIMIPAROUS MOTHERS AGED BETWEEN 20 AND 34 IN PUBLIC ACCOMMODATION, QUEENSLAND, 1998, ONSET OF LABOUR BY METHOD OF DELIVERY**

Onset of labour	Method of delivery									
	Spontaneous Vertex		Forceps/Vacuum		Caesarean Section		Other		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%
Spontaneous - not augmented	2,231	41.2	275	22.9	187	15.5	-	-	2,693	34.4
Spontaneous - augmented	1,929	35.6	547	45.5	425	35.2	2	66.7	2,903	37.1
Induced	1,254	23.2	380	31.6	475	39.4	1	33.3	2,110	27.0
No labour	-	-	-	-	120	9.9	-	-	120	1.5
<b>Total</b>	<b>5,414</b>	<b>100.0</b>	<b>1,202</b>	<b>100.0</b>	<b>1,207</b>	<b>100.0</b>	<b>3</b>	<b>100.0</b>	<b>7,826</b>	<b>100.0</b>


(a) Includes babies born between 37 and 41 completed gestational weeks.



**TABLE 6.12 MOTHERS WITH A PREVIOUS CAESAREAN SECTION, QUEENSLAND, 1998**  
**METHOD OF DELIVERY BY ONSET OF LABOUR <sup>(a)</sup>**

Onset of labour	Method of delivery									
	Spontaneous Vertex		Forceps/ Vacuum		Caesarean Section		Other		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%
Spontaneous - not augmented	580	45.4	65	32.5	632	14.7	10	58.8	1,287	22.2
Spontaneous - augmented	469	36.7	88	44.0	137	3.2	1	5.9	695	12.0
Induced	229	17.9	47	23.5	105	2.4	6	35.3	387	6.7
No labour	-	-	-	-	3,423	79.7	-	-	3,423	59.1
<b>Total</b>	<b>1,278</b>	<b>100.0</b>	<b>200</b>	<b>100.0</b>	<b>4,297</b>	<b>100.0</b>	<b>17</b>	<b>100.0</b>	<b>5,792</b>	<b>100.0</b>

(a) In the case of multiple births data from the first birth are used.




**TABLE 6.13 MOTHERS DELIVERING IN QUEENSLAND, 1998  
SELECTED LABOUR AND DELIVERY COMPLICATIONS**

Complication <sup>(a)</sup>	Mothers	
	No.	%
Labour < 4 hours	9,723	20.5
Fetal distress	4,285	9.0
Uterine inertia	3,289	6.9
Prolonged second stage	2,300	4.8
Cord entanglement	1,215	2.6
Primary post-partum haemorrhage	1,165	2.5
Obstructed labour	881	1.9
Retained placenta or membranes (without haemorrhage)	848	1.8
Maternal distress	69	0.1
Other	20,155	42.5
<b>Total mothers</b>	<b>47,450</b>	

(a) Reporting of multiple complications was permitted, so numbers cannot be cumulated.

Labour and delivery complications were reported for 31,559 (66.5 per cent) of mothers.




**TABLE 6.14 BABIES BORN AFTER A PERIOD OF LABOUR, QUEENSLAND, 1998  
METHOD OF DELIVERY BY SELECTED LABOUR AND DELIVERY COMPLICATIONS**

Selected complication <sup>(a)</sup>	Method of delivery									
	Spontaneous		Forceps/ Vacuum		Caesarean		Other		Total <sup>(b)</sup>	
	Vertex				Section					
	No.	%	No.	%	No.	%	No.	%	No.	%
Fetal distress	1,378	4.3	1,185	27.7	1,572	32.8	23	5.5	4,158	10.0
Meconium liquor	4,925	15.3	808	18.9	955	19.9	49	11.8	6,737	16.1
Long labour	784	2.4	1,819	42.5	325	6.8	9	2.2	2,937	7.0
Primary post-partum haemorrhage	870	2.7	172	4.0	57	1.2	14	3.4	1,113	2.7
Uterine inertia	205	0.6	836	19.5	2,270	47.3	5	1.2	3,316	7.9
Obstructed labour	442	1.4	191	4.5	244	5.1	5	1.2	882	2.1
Lacerations/tears to perineum/vulva	15,335	47.5	1,516	35.4	41	0.9	75	18.1	16,967	40.6
<b>Total babies born after a period of labour</b>	<b>32,279</b>		<b>4,278</b>		<b>4,797</b>		<b>415</b>		<b>41,769</b>	

(a) Reporting of multiple conditions permitted, so numbers cannot be cumulated.

(b) Total includes one baby for whom the method of delivery was not stated.




**TABLE 6.15      MOTHERS DELIVERING IN QUEENSLAND, 1998**  
**SELECTED PROCEDURES AND OPERATIONS PERFORMED**  
**DURING PREGNANCY, LABOUR AND THE PUERPERIUM**

Procedures and Operations <sup>(a)</sup>	Mothers	
	No.	%
Obstetric ultrasound	46,670	98.4
Monitoring of fetal heart during labour	24,858	52.4
Amniocentesis	1,907	4.0
Removal of retained placenta	776	1.6
Encirclement suture of cervix	116	0.2
X-ray pelvimetry	10	0.0
Other	1,055	2.2
<b>Total mothers</b>	<b>47,450</b>	

(a) Reporting of multiple procedures was permitted so numbers cannot be cumulated.

Procedures and operations performed during pregnancy, labour and the puerperium were reported for 47,012 (99.1 per cent) mothers. The marked increase in reporting of procedures and operations is most likely attributable to improved reporting of obstetric ultrasound.




**TABLE 6.16 MOTHERS DELIVERING IN QUEENSLAND, 1998  
SELECTED PUERPERIUM COMPLICATIONS**

Complication <sup>(a)</sup>	Mothers	
	No.	%
Complications of obstetrical surgical wounds	369	0.8
Infection of genitourinary tract	299	0.6
Puerperial pyrexia	153	0.3
Post-partum haemorrhage	196	0.4
Infection of breast	122	0.3
Disorders of the breast excluding infection	239	0.5
Venous complications	107	0.2
Anaemia	161	0.3
Other	1,240	2.6
<b>Total mothers</b>	<b>47,450</b>	

(a) Reporting of multiple complications was permitted so numbers cannot be cumulated.

At least one puerperium complication was reported for 2,714 (5.7 per cent) of mothers.

The scope of the Queensland Perinatal Data Collection ceases at the point of formal separation - that is discharge, transfer, or death. Whilst the average length of stay has continued to reduce in recent years, the proportion of mothers with at least one puerperium complication has gradually increased. It is suspected that this may be an indication of improved reporting.



**TABLE 6.17    MOTHERS DELIVERING VAGINALLY, QUEENSLAND, 1998**  
**PERINEAL STATUS BY NUMBER OF PREVIOUS DELIVERIES**

Perineal status	Number of previous deliveries							
	0		1 - 4		5 or more		Total <sup>(a)</sup>	
	No.	%	No.	%	No.	%	No.	%
Intact	4,050	29.0	11,420	52.0	589	78.4	16,060	43.8
<i>Without other lower genital tract trauma</i>	2,952	21.1	10,476	47.7	566	75.4	13,995	38.2
Lacerated sutured	5,085	36.4	6,448	29.4	81	10.8	11,614	31.7
Lacerated unsutured	875	6.3	2,085	9.5	67	8.9	3,027	8.3
Episiotomy	3,954	28.3	1,987	9.1	13	1.7	5,954	16.2
Not stated	-	-	1	0.0	1	0.1	2	0.0
<b>Total</b>	<b>13,964</b>	<b>100.0</b>	<b>21,941</b>	<b>100.0</b>	<b>751</b>	<b>100.0</b>	<b>36,657</b>	<b>100.0</b>

(a) Includes not stated number of previous confinements.

**TABLE 6.18      MOTHERS DELIVERING VAGINALLY, QUEENSLAND, 1998**  
**PERINEAL STATUS BY SELECTED GENITAL TRACT TRAUMA**

Selected genital tract trauma <sup>(a)</sup>	Perineal status									
	Intact		Lacerated- sutured		Lacerated- unsutured		Episiotomy		Total <sup>(b)</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%
First degree perineal laceration	-	-	1,194	10.3	623	20.6	27	0.5	1,844	5.0
Second degree perineal laceration	-	-	1,396	12.0	49	1.6	67	1.1	1,512	4.1
Third degree perineal laceration	-	-	108	0.9	2	0.1	101	1.7	211	0.6
Fourth degree perineal laceration	-	-	7	0.1	-	-	8	0.1	15	0.0
Unspecified perineal laceration	45	0.3	2,997	25.8	749	24.7	216	3.6	4,007	10.9
Other trauma to perineum or vulva	1,660	10.3	222	1.9	185	6.1	51	0.9	2,118	5.8
High vaginal laceration	395	2.5	129	1.1	17	0.6	37	0.6	578	1.6
<b>Mothers delivering vaginally</b>	<b>16,060</b>		<b>11,614</b>		<b>3,027</b>		<b>5,954</b>		<b>36,657</b>	

(a) Reporting of multiple genital tract conditions permitted so numbers cannot be cumulated.

In the case of multiple births details from the first birth are used.

(b) Includes not stated perineal status.

**TABLE 6.19 MOTHERS AGED BETWEEN 20 AND 34 IN PUBLIC ACCOMMODATION WITH A SPONTANEOUS VERTEX DELIVERY OF A SINGLETON, TERM <sup>(a)</sup> BABY QUEENSLAND, 1998, PERINEAL STATUS BY ACCOUCHEUR**

Perineal Status	Accoucheur												Total <sup>(b)</sup>	
	Other													
	Obstetrician		Medical officer		Midwife		Student midwife		Medical student		Other		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b><i>Primiparous mothers</i></b>														
Intact	17	23.6	100	26.0	1,448	35.9	226	31.8	69	33.7	5	62.5	1,866	34.5
<i>Without other lower genital tract trauma</i>	15	20.8	74	19.3	993	24.6	153	21.5	52	25.4	2	25.0	1,289	23.8
Lacerated sutured	28	38.9	138	35.9	1,703	42.2	322	45.4	87	42.4	3	37.5	2,281	42.1
Lacerated unsutured	3	4.2	21	5.5	377	9.3	69	9.7	14	6.8	-	-	484	8.9
Episiotomy	24	33.3	125	32.6	506	12.5	93	13.1	35	17.1	-	-	783	14.5
<b>Total</b>	<b>72</b>	<b>100.0</b>	<b>384</b>	<b>100.0</b>	<b>4,034</b>	<b>100.0</b>	<b>710</b>	<b>100.0</b>	<b>205</b>	<b>100.0</b>	<b>8</b>	<b>100.0</b>	<b>5,414</b>	<b>100.0</b>
<b><i>Multiparous mothers</i></b>														
Intact	44	43.1	385	57.2	5,530	60.8	1,033	60.2	282	55.0	89	67.9	7,364	60.2
<i>Without other lower genital tract trauma</i>	40	39.2	361	53.6	5,030	55.3	930	54.2	261	50.9	82	62.6	6,705	54.8
Lacerated sutured	28	27.5	169	25.1	2,117	23.3	446	26.0	143	27.9	26	19.8	2,929	23.9
Lacerated unsutured	15	14.7	60	8.9	1,077	11.8	179	10.4	70	13.6	16	12.2	1,418	11.6
Episiotomy	15	14.7	59	8.8	370	4.1	57	3.3	18	3.5	-	-	519	4.2
<b>Total</b>	<b>102</b>	<b>100.0</b>	<b>673</b>	<b>100.0</b>	<b>9,094</b>	<b>100.0</b>	<b>1,715</b>	<b>100.0</b>	<b>513</b>	<b>100.0</b>	<b>131</b>	<b>100.0</b>	<b>12,230</b>	<b>100.0</b>

(a) Includes babies born between 37 and 41 completed gestational weeks.

(b) Includes 3 mothers with a not stated accoucheur.