



Transjugular Liver Biopsy

Facility:

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

D. Risks of Iodinated Contrast for patients with renal impairment

Specific Risks of Iodinated Contrast to patient's identified as having *Renal Impairment*.

- Giving the Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).

E. Patient consent

I acknowledge that the doctor/doctor delegate has explained the proposed procedure.

I understand;

- the risks and complications, including the risks that are specific to me.
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor/doctor delegate or my Acute Resuscitation Plan.
- a doctor/doctor delegate undergoing further training may conduct this procedure.

I have been given the following Patient Information Sheet/s:

- Transjugular Liver Biopsy**
- Iodinated Contrast**
- About Your Anaesthetic** (if required)

- I was able to ask questions and raise concerns with the doctor/doctor delegate about the proposed procedure and its risks. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time including after I have signed this form but, preferably following a discussion with my doctor/doctor delegate.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.
- I understand that Queensland Health may release my relevant de-identified information obtained from this and related procedures for education and training of health professionals.

On the basis of the above statements,

I request to have the procedure

Name of Patient:

Signature:

Date:

Patients who lack capacity to provide consent

Consent must be obtained from a substitute decision maker/s in the order below.

Does the patient have an Advance Health Directive (AHD)?

Yes ► Location of the original or certified copy of the AHD:

No ► Name of Substitute Decision Maker/s:
Signature:
Relationship to patient:
Date: PH No:

Source of decision making authority (tick one):

- Tribunal-appointed Guardian
- Attorney/s for health matters under Enduring Power of Attorney or AHD
- Statutory Health Attorney
- If none of these, the Adult Guardian has provided consent. Ph 1300 QLD OAG (753 624)

F. Doctor/delegate Statement

I have explained to the patient all the above points under:

- the Patient Consent section (E)
- Iodinated Contrast - Patients with Renal Impairment Section (D) (for renal impaired patients only)

and I am of the opinion that the patient/substitute decision-maker has understood the information.

Name of Doctor/delegate:
Designation:

Signature: **Date:**

G. Interpreter's statement

I have given a sight translation in

.....
(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

Name of Interpreter:

Signature: **Date:**

DO NOT WRITE IN THIS BINDING MARGIN

Consent Information - Patient Copy

Transjugular Liver Biopsy

1. What is a Transjugular Liver Biopsy?

A transjugular liver biopsy is where a sample of tissue is taken from your liver. This is done by placing a catheter (hollow tube) and needle into your jugular vein which is guided into your liver. The sample is taken and sent to pathology for testing.

Iodinated 'Contrast' (once called x-ray dye) is used to map your veins so the doctor can place the catheter into the hepatic vein. For more information on Iodinated Contrast and the risks involved in its use, please read the **Iodinated Contrast Patient Information Sheet**. (*If you do not have this information sheet please ask for one*).

This procedure is performed in medical imaging with guidance from imaging machines such as ultrasound and x-ray.

2. Will there be any discomfort, is any anaesthetic needed?

This procedure will require an injection of local anaesthetic and the use of a sedation or general anaesthetic. If you are booked for a general anaesthetic please read the **About your Anaesthetic** patient information sheet (*If you do not have this information sheet please ask for one*).

3. What is sedation?

Sedation is the use of drugs that give you a 'sleepy-like' feeling. It makes you feel very relaxed during a procedure that may be otherwise unpleasant or painful. You may remember some or little about what has occurred during the procedure.

This procedure may only have a light sedation. You need to be able to fully co-operate at times by holding your breath when instructed by the doctor.

Sedation is generally very safe but has a risk with side effects and complications. Whilst these are usually temporary, some of them may cause long-term problems.

The risk to you will depend on:

- whether you have any other illness
- personal factors, such as whether you smoke or are overweight.

4. Preparation for the procedure

The medical imaging department will give you instructions on how to prepare for your procedure.

- You will be told when to have your last meal and drink. This is to make sure your stomach is empty so that if you vomit during the procedure there will be nothing to go into your lungs.
- Please tell the staff if you are or suspect you might be pregnant.
- *If you take Aspirin, Warfarin, Clopidogrel (Plavix and Iscover) or Dipyridamole (Persantin and Asasantin) or any other drug that is used to thin*

your blood ask your doctor/health practitioner if you should stop taking it before the procedure as it may affect your blood clotting.

- *List or bring all your prescribed drugs, those drugs you buy over the counter, herbal remedies and supplements.*
- *Do not drink any alcohol* and *stop recreational drugs* 24 hours before the procedure as this may alter the affects of the sedation anaesthetic. If you have a drug habit please tell your doctor.

5. During the procedure

A fine needle (IV cannula) will be inserted into a vein in your arm.

The Radiologist (x-ray doctor) will inject local anaesthetic into the skin on your neck.

A small cut will be made.

Using ultrasound as a guide the Radiologist will insert a needle through the cut and into your jugular vein.

You must remain as still as possible. At times, you may be asked to hold your breath.

The catheter will be inserted into the vein and the needle removed. Contrast will be injected as x-ray pictures are taken.

Once the catheter is in place, a long needle is then passed through the catheter to obtain the tissue sample.

It is not unusual for this step to be repeated.

At the end of the procedure, the needle and catheter will be removed and pressure will be applied over the area where the catheter went into your skin (puncture site). A dressing will be applied.

6. After the procedure

The recovery time varies depending on the sedation that has been given. It varies between 4 hours to 6 hours.

The IV cannula will be removed after you have recovered.

Staff will discuss with you what level of activity is suitable after your procedure.

7. What are the risks of this specific procedure?

The risks and complications with this procedure can include but are not limited to the following.

Common risks and complications include:

- Minor pain, bruising and/or infection from the IV cannula. This may require treatment with antibiotics.
- Pain or discomfort at the puncture site. This may require medication.
- Bleeding or bruising may occur. This is more common if you take Aspirin, Warfarin, Clopidogrel

1. What is Iodinated Contrast?

The medical imaging procedure your doctor has asked you to have uses Iodinated 'Contrast' (once called X-ray dye). Contrast is a colourless liquid which includes iodine. Contrast is injected into your blood stream to allow your organs to be seen more clearly on x-rays. Contrast is not a dye. It does not stain the inside of your body. Your doctor needs to use Contrast to be able to get all the information needed to assist with your diagnosis.

This information sheet must be read together with the information sheet of the procedure you are booked for (if you do not have this information sheet please ask for one).

2. During the procedure

When the Contrast is injected you may feel:

- A very warm or 'flushed' feeling over your body, this may also make you think you have passed urine. You will *not* pass urine – it is only a feeling.
- A 'metallic' taste or smell may also happen. This usually lasts less than a minute.

3. After the procedure

It is recommended that you drink 2 to 4 glasses of water after your procedure to help flush the Contrast from your body.

Contrast does not affect your ability to carry out normal activities; you should be able to continue with your day as normal.

4. Precautions

Contrast is not suitable for some people; you will be asked a series of questions before it is given to you. Your answers allow staff to identify any risk factors that you may have.

- Please tell the staff if you are or suspect you might be pregnant or are breastfeeding.

Kidney function

- Contrast is removed from your blood by your kidneys through your urine. It is easily removed from the body of people who have normal kidney function.
- Giving Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).
- You may be asked to have a simple blood test to find out the level of their kidney function.

Diabetic Drug interactions - Metformin

(Other Drug names: Avandamet, Diabex, Diaformin, Formet, Glucohexal, Glucomet, Glucophage, Glucovance, Metforbell)

If kidneys suffer damage from the Contrast then the kidneys may not be able to remove Metformin from the body. It is safer to briefly stop taking Metformin when having Contrast. Staff will inform you when to stop and when it is safe to take Metformin again.

Contact your GP to monitor your diabetes if you are told to stop your Metformin.

5. What are the risks of Iodinated Contrast?

The risks and complications with this injection can include but are not limited to the following.

Common risks and complications include:

- No known common risks.

Less common risks and complications include:

- Injected Contrast may leak outside of the blood vessel, under the skin and into the tissue. This may require treatment. In very rare cases, further surgery could be required if the skin breaks down.
- Acute Renal Failure occurs when one or both of your kidneys suddenly stop working. Failure can last for days or weeks. It may take the kidneys a long time to regain their previous level of function and you may require dialysis to filter your blood during this time. There is a risk your kidneys could be permanently damaged. To reduce this risk the smallest possible dose of Contrast will be given.
- The injection may not be possible due to medical and/or technical reasons.

Rare risks and complications include:

- Allergic reactions occur within the first hour with most happening in the first 5 minutes. Late reactions have been known to occur up to a week after the injection.

Note: Allergy to topical iodine and/or seafood does not imply an allergy to Iodinated Contrast.

The reactions vary from:

Mild – hives, sweating, sneezing, coughing, nausea.

Moderate – wide spread hives, headaches, facial swelling, vomiting, shortness of breath.

Severe – Severe reactions are rare but include: life-threatening heart palpitations, very low blood pressure, throat swelling, fits and/or cardiac arrest.

- Death as a result of Iodinated Contrast is *very* rare.

6. What are the safety issues when you leave the hospital?

Go to your nearest Emergency Department or GP if you become unwell.