



SW1106

v2.00 - 05/2022

DO NOT WRITE IN THIS BINDING MARGIN

© State of Queensland (Queensland Health) 2022
Licensed under: https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en
For further information contact Queensland Clinical Guidelines, email: guidelines@health.qld.gov.au
For permissions beyond the scope of this licence contact:
Intellectual Property Officer email: ip_officer@health.qld.gov.au, phone: (07) 3234 1479.



Nomogram: Jaundice Management
For baby greater than 38 weeks gestation

Facility:

Comments:
Time of birth (24hr):
DCT:
Baby's blood group:
Mother's blood group:

(Affix identification label here)
URN:
Family name:
Given name(s):
Address:
Date of birth:
Sex: M F I

- 1. In the presence of risk factors (sepsis, haemolysis, acidosis or asphyxia) use the lower line.
2. If baby is greater than 12 hours old with total serum bilirubin (TSB) 1-50 micromol/L below the line, repeat the TSB within 6-24 hours.
3. Babies under phototherapy:
a. Consider measuring the TSB 4-6 hourly until the rise of serum bilirubin is known to be controlled, then measure TSB 12-24 hourly.
b. If the TSB is greater than 50 micromol/L below line, stop phototherapy and recheck in 12-24 hours.
4. If baby presents with TSB above threshold and the TSB is not expected to be below the threshold after 6 hours of intensive phototherapy, an exchange transfusion is indicated.
5. If there are signs of bilirubin encephalopathy an immediate exchange transfusion is recommended.

Baby greater than 38 weeks gestation




Table with 5 columns: Date, Time (24hr), TSB Result, Start time (24hr), End time (24hr). Multiple rows for recording phototherapy sessions, each with a 'Comments:' field.