



**Queensland  
Government**

## Iron Infusion Consent

Adult (18 years and over) *OR*  
 Child/Young Person (under 18 years)

Facility: .....

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex:  M  F  I

### A. Does the patient have capacity?

#### Complete for ADULT patient only

- Yes → **GO TO section B**  
 No → **COMPLETE section A**

You must adhere to the Advance Health Directive (AHD), or if there is no AHD, the consent obtained from a substitute decision-maker in the following order: Category 1. Tribunal-appointed guardian; 2. Enduring Power of Attorney; or 3. Statutory Health Attorney.

Name of substitute decision-maker:

Category of substitute decision-maker:

#### Complete for CHILD/YOUNG PERSON patient only

- Yes Although the patient is a child/young person, the patient may be capable of giving informed consent and having sufficient maturity, understanding and intelligence to enable them to fully understand the nature, consequences and risks of the proposed procedure/treatment/investigation/examination and the consequences of non-treatment – ‘Gillick competence’ (*Gillick v West Norfolk and Wisbech Area Health Authority* [1986] AC 112)  
 → **GO TO section B**
- No Parent/legal guardian/other person\* with parental rights and responsibilities to provide consent and complete this form  
 → **COMPLETE section A**

\*Formal arrangements, such as parenting/custody orders, adoption, or other formally recognised carer/guardianship arrangements. Refer to the Queensland Health ‘Guide to Informed Decision-making in Health Care’ and local policy and procedures. Complete the source of decision-making authority as applicable below.

If applicable, source of decision-making authority (*tick one*):

- Court order →  Court order verified  
 Legal guardian →  Documentation verified  
 Other person\* →  Documentation verified

Name of parent/legal guardian/other person\*:

Relationship to child/young person:

### B. Is an interpreter required?

If yes, the interpreter has:

- provided a sight translation of the informed consent form in person  
 translated the informed consent form over the telephone

Name of interpreter:

Interpreter code:

Language:

### C. Patient *OR* substitute decision-maker *OR* parent/legal guardian/other person\* requests the following procedure(s)

Iron infusion

- Ferric Carboxymaltose (Ferinject®)  
 Ferric Derisomaltose (Monofer®)  
 Iron Polymaltose (Ferrosig®)  
 Iron Sucrose (Venofer®)

Is this off-label use?  Yes  No

### D. Risks specific to the patient in having an iron infusion

(Doctor/clinician to document additional risks not included in the patient information sheet):

### E. Risks specific to the patient in *not* having an iron infusion

(Doctor/clinician to document specific risks in not having an iron infusion):

### F. Alternative treatment options

(Doctor/clinician to document alternative treatment not included in the patient information sheet):

### G. Information for the doctor/clinician

The information in this consent form is not intended to be a substitute for direct communication between the doctor/clinician and the patient *OR* substitute decision-maker *OR* parent/legal guardian/other person\*.

I have explained to the patient *OR* substitute decision-maker *OR* parent/legal guardian/other person\* the contents of this form and am of the opinion that the information has been understood.

Name of doctor/clinician:

Designation:

Signature:

Date:

DO NOT WRITE IN THIS BINDING MARGIN

v1.00  
 Clinical content review: 2021  
 Clinical check: 08/2021  
 Published: 09/2021



SW9511

IRON INFUSION CONSENT



**Queensland  
Government**

## Iron Infusion Consent

Adult (18 years and over) *OR*  
Child/Young Person (under 18 years)

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex:  M  F  I

### H. Patient *OR* substitute decision-maker *OR* parent/ legal guardian/other person\* consent

I acknowledge that the doctor/clinician has explained:

- the “Iron infusion” patient information sheet
- the medical condition and proposed treatment, including the possibility of additional treatment
- the general and specific risks and benefits of the procedure
- the prognosis, and risks of not having the procedure
- alternative treatment options
- that there is no guarantee the procedure will improve the medical condition
- that if a life-threatening event occurs during the procedure:
  - an adult/young person will be treated based on documented discussions (e.g. AHD or ARP [Acute Resuscitation Plan])
  - a child’s health care will be provided in accordance with good clinical practice and in the best interests of the patient
- that a doctor/clinician other than the consultant/specialist may assist with/conduct the clinically appropriate treatment; this may include a doctor/clinician undergoing further training under supervision
- that if the doctor/clinician wishes to record video, audio or images during the procedure where the recording is not required as part of the treatment (e.g. for training or research purposes), I will be asked to sign a separate consent form. If I choose not to consent, it will not adversely affect my access, outcome or rights to medical treatment in any way.

I was able to ask questions and raise concerns with the doctor/clinician.

I understand I have the right to change my mind regarding consent at any time, including after signing this form (*this should be in consultation with the doctor/clinician*).

**I/substitute decision-maker/parent/legal guardian/other person\* have received the following consent and patient information sheet(s):**

“Iron infusion”

On the basis of the above statements,

**I/substitute decision-maker/parent/legal guardian/other person\* consent to having an iron infusion.**

Name of patient/substitute decision-maker/parent/legal guardian/other person\*:

Signature:

Date:

If the patient is a young person:

I am not aware of any legal or other reason that prevents me from providing unrestricted consent for this young person for this treatment (*not applicable if the young person is Gillick competent and signs this form*).

**A copy of this form should be given to the patient/substitute decision-maker/parent/legal guardian/other person\* to read carefully and allow time to ask any questions about the procedure. The consent form and patient information sheet should be included in the patient's medical record.**



## 1. What is an iron infusion and how will it help me/the patient?

Our bodies need iron. Iron is used to make haemoglobin, this is the part of the red blood cell that transports oxygen around the body. Iron is also important for muscle strength, energy and mental function. If your iron levels are low you may feel tired and not be able to perform your normal daily activities. As the iron levels drop lower, the haemoglobin levels drop below normal. This is known as iron deficiency anaemia.

The most common way to treat iron deficiency is to take iron by mouth as a tablet or liquid. This works for most people and is usually tried first.

You might need an iron infusion if you:

- are not able to take iron tablets/liquids
- are not responding to iron tablets/liquids
- need to increase your iron levels quickly (e.g. before major surgery, late in pregnancy or to avoid a blood transfusion)
- if you have chronic kidney disease, chronic heart failure, or inflammatory bowel disease.

Sometimes more than one iron infusion is required to fully elevate your iron levels.

An intravenous (IV) iron infusion is an infusion made up of iron, not blood, that is infused directly into your bloodstream. A plastic cannula is placed into one of your veins and is attached to a pump that infuses the iron into your body. Iron infusions work faster to replace iron than tablets or liquids taken by mouth.

Staining of the skin from intravenous iron is uncommon, but when it occurs **the stain can be long lasting or permanent** (see *image above*). The leakage occurs in the area around the cannula (drip site) into the surrounding tissues.



*Image: Iron staining.*

*Source: Canning, M & Grannell, L. 2020, 'A Stain on iron therapy,' Australian Prescriber, vol. 43, no. 5, pp.160-3, retrieved 29/08/2021 ([www.nps.org.au/assets/p160-Canning-Grannell.pdf](http://www.nps.org.au/assets/p160-Canning-Grannell.pdf)). Image used under CC BY-NC-ND 4.0 licence.*

## Iron use in children

The product information for each of the following products state they are not indicated for use in children; however, iron infusions are commonly used in children. This is called off-label use and is common in paediatric medical practice. If you have any questions about off-label use, please discuss these with your doctor/nurse/midwife. For further information on off-label use in children go to section 6 "*Where can I find support or more information?*". The information in these guides refers to the best available evidence and best practice protocols that support the safety and benefit of these products for paediatric use.

All the iron products listed on this consent form have Australian marketing approval from the Therapeutic Goods Administration (TGA). The Australian approved product information lists the approved indications (uses) and any age restrictions. Off-label use occurs when an iron infusion is prescribed outside the approved product information (either the indication or the age restriction) and is common in paediatric medical practice. A doctor/nurse/midwife will discuss the suitable treatment options for your child. The indications and age restrictions for iron infusion are as follows:

- **Ferric Carboxymaltose (Ferinject®)**
  - Treatment of iron deficiency when oral iron preparations are ineffective or cannot be used. The diagnosis must be based on laboratory tests. Precaution: The product information states that Ferric Carboxymaltose is not recommended in children under 14 years.
- **Ferric Derisomaltose (Monofer®)** – Product information states that Ferric Derisomaltose Iron Supplement is for Adults 18 years and older, for use in iron deficiency when oral preparations are ineffective, inappropriate or when rapid iron delivery is necessary.
- **Iron Polymaltose (Ferrosig®)** – Product information states that there is no age restriction. Treatment of iron deficiency anaemia when oral therapy is contraindicated, gastrointestinal absorption of iron is defective or when patient noncompliance or persistent gastrointestinal intolerance makes oral therapy impractical. Iron supplement for Iron deficiency anaemia where oral treatment not appropriate or gastrointestinal absorption is defective.
- **Iron Sucrose (Venofer®)** – Treatment of iron deficiency anaemia in patients undergoing long-term haemodialysis and who are receiving supplemental erythropoietin therapy (a drug used to stimulate red blood cell production in bone marrow for anaemia). The diagnosis of iron deficiency must be based on appropriate laboratory tests. Precaution: The product information states that the safety and efficacy of Iron Sucrose in children has not been established.

## Preparing for the treatment

Before you have an iron infusion, you must tell your doctor/nurse/midwife and the nurses performing the infusion if you:

- are pregnant/trying to get pregnant, as an IV iron infusion should be avoided in the first trimester
- have a history of asthma, eczema or other allergies
- have had a reaction to any type of iron injection or infusion in the past
- have a history of high iron levels (haemochromatosis) or liver problems
- have (or may have) an infection at the moment.

You may want to ask your doctor/nurse/midwife about the following:

- Why do I need an IV iron infusion?
- What are the other options?
- How many infusions will I need to get enough iron?
- *(If you are currently taking iron tablets/liquids)* When do I stop taking the iron tablets/liquids and will I need to use them again?
- How long will it take for my iron levels to improve?
- Any questions about side effects that may worry you.

## The day of the treatment

There is preparation needed for an iron infusion. Here are some useful tips:

- It is helpful if you are well hydrated prior to your iron infusion so that putting in a cannula may be easier.
- Have your regular breakfast/lunch – you do not need to fast.
- Take all your regular medications.
- Tell your nurse/midwife administering the iron infusion if you are pregnant or not.
- If you experience any side effects, inform your nurse/midwife immediately.
- Adults can drive home after the iron infusion and all patients can resume usual activities (unless you experienced an unexpected reaction and medical staff inform you otherwise).
- Very little iron crosses into breast milk so breast feeding mothers may safely breastfeed.

## During the treatment

The duration of the iron infusion varies depending on the product used. Your iron infusion may take 15 to 30 minutes, 1 to 2 hours or 4 to 6 hours.

You will also need to allow time for the IV cannula to be inserted and removed, and for clinical observation before and after the iron infusion (30 to 60 minutes).



## 2. What are the risks?

There are risks and complications with this procedure. There may also be risks specific to each person's individual condition and circumstances. Please discuss these with the doctor/nurse/midwife and ensure they are written on the consent form before you sign it. Risks include but are not limited to the following:

### Common risks and complications

- low blood pressure
- palpitations
- facial flushing
- dizziness
- vomiting and nausea
- headache
- injection site reactions (*see also below skin staining in rare risks*)
- joint and/or muscle pain
- changes in taste (e.g. metallic)
- (*Ferric carboxymaltose [Ferinject®] only*) hypophosphataemia (low blood phosphate level). This can cause muscle weakness, respiratory failure, heart failure and in rare circumstances osteomalacia (bone softening), fractures, seizures and/or coma. You may require oral medication to raise your blood phosphate level.

### Uncommon risks and complications

- dizziness and feeling faint
- chest and/or back pain
- chills and fever
- skin irritation and rash
- swelling of the face, mouth and limbs
- generalised lymphadenopathy (swollen lymph nodes)
- difficult breathing.

## Rare risks and complications

- anaphylaxis – rare and usually occurs in the first few minutes of the iron infusion and is characterised by breathing difficulties, which may be life threatening. You will be closely monitored whilst the iron infusion is given and for at least 30 minutes
- rapid heart beat (tachycardia) and low blood pressure
- skin staining (brown discoloration) may occur due to leakage of iron into the tissue around the cannula (drip) site. This is not common, but the stain can be long lasting or permanent (*see image*)
- inform the nurse straight away of any pain, discomfort, burning, prickling, redness, staining or swelling at the cannula (drip) site
- death as a result of this procedure is possible. You will be closely monitored whilst the iron infusion is given and for at least 30 minutes following completion.

This treatment does not require an anaesthetic.

## What are the risks of not having an iron infusion?

There may be health consequences if you choose not to have the proposed treatment. Please discuss these with the doctor/nurse/midwife.

If you choose not to have the treatment, you will not be required to sign a consent form.

If you have signed a consent form, you have the right to change your mind at any time prior to the treatment. Please contact the doctor/nurse/midwife to discuss.



## 3. Are there alternatives?

Making the decision to have a procedure requires the patient/substitute decision-maker/parent/legal guardian/other person\* to understand the options available. Please discuss any alternative treatment options with your doctor/nurse/midwife before signing the consent form.



## 4. What should I expect after the treatment?

In some cases, iron tablets/liquids are recommended after the infusion. If oral iron therapy is recommended it should not start until at least one week after your IV iron infusion as the iron within the tablets/liquids will not be absorbed.

**Tell your doctor/nurse/midwife or seek urgent medical attention/call an ambulance (000) if you have:**

- chest pain
- trouble breathing
- dizziness
- neck/mouth swelling.



## 5. Who will be performing the treatment?

A doctor/nurse/midwife other than the consultant/specialist may assist with/conduct the clinically appropriate treatment. This could be a doctor/nurse/midwife undergoing further training, all trainees are supervised according to relevant professional guidelines.

If you have any concerns about which doctor/nurse/midwife will be performing the treatment, please discuss with the doctor/nurse/midwife.



## 6. Where can I find support or more information?

Hospital care: before, during and after is available on the Queensland Health website [www.qld.gov.au/health/services/hospital-care/before-after](http://www.qld.gov.au/health/services/hospital-care/before-after) where you can read about your healthcare rights.

More information on iron infusions can be found on the TGA website:

[www.tga.gov.au/sites/default/files/auspar-ferric-carboxymaltose-191001.pdf](http://www.tga.gov.au/sites/default/files/auspar-ferric-carboxymaltose-191001.pdf)  
[www.tga.gov.au/sites/default/files/auspar-ferric\\_derisomaltose-181105.pdf](http://www.tga.gov.au/sites/default/files/auspar-ferric_derisomaltose-181105.pdf)

More information on off-label use can be found at:

[www.blood.gov.au/system/files/Corrected-Paed-and-Neonatal-IDA-Guide-dosing-for-iron-polymaltose.pdf](http://www.blood.gov.au/system/files/Corrected-Paed-and-Neonatal-IDA-Guide-dosing-for-iron-polymaltose.pdf)

[www.mja.com.au/journal/2006/185/10/label-use-medicines-consensus-recommendations-evaluating-appropriateness](http://www.mja.com.au/journal/2006/185/10/label-use-medicines-consensus-recommendations-evaluating-appropriateness)

Staff are available to support patients' cultural and spiritual needs. If you would like cultural or spiritual support, please discuss with your doctor/nurse/midwife.

Queensland Health recognises that Aboriginal and Torres Strait Islander patients will experience the best clinical care when their culture is included during shared decision-making.



## 7. Questions

Please ask the doctor/nurse/midwife if you do not understand any aspect of this patient information sheet or if you have any questions about your/the patient's medical condition, treatment options and proposed treatment.

If you require additional information about the intravenous iron preparation prescribed for you, please speak to the clinician/doctor/nurse/pharmacist who can arrange to provide you with a Consumer Medicines Information (CMI) leaflet.



## 8. Contact us

**In an emergency, call Triple Zero (000).**

If it is not an emergency, but you have concerns, contact 13 HEALTH (13 43 25 84), 24 hours a day, 7 days a week.