


Queensland Community Pharmacy Scope of Practice Pilot

Acute Otitis Media – Clinical Practice Guideline

Guideline Overview





'Red flag' warning signs at patient presentation that necessitate:

a. Urgent referral to a medical practitioner including emergency care (without concurrent antibiotic therapy):

- Recent trauma to the ear or head (this may be associated with hearing loss or balance issues)
- Severe symptoms or rapidly worsening symptoms including intense pain, headache or facial paralysis/palsy, red, swollen and tender behind the ear (possible acute mastoiditis)
- Tympanic membrane (TM) 'red flag' signs:
 - Crust/scab or granulation in attic region (discharge/odour may be present)
 - Severely retracted (sucked in) TM or retraction pocket in attic area of eardrum
 - Dull white mass behind TM
 - Perforation in attic region, near the edge of the TM or medium to large in size (refer to Figure 1).
- Signs or symptoms of sepsis including fever or hypothermia, significant concern from parent/carer, respiratory distress, vomiting, tachycardia, rash, hypotension and/or cold extremities or other significant systemic symptoms including weakness, irritability, difficulty sleeping and/or loss of appetite.

b. Antibiotic therapy may be provided concurrently to medical practitioner referral:

- Otic discharge/otorrhea present
- Children with symptoms that worsen or do not improve after 48 hours of regular analgesia and supportive management
- Symptoms are recurrent (3 or more episodes in 6 months or 4 or more episodes in 12 months)
- The patient has persistent otitis media with effusion (glue ear)
- Children with signs and symptoms of a complication, or at high risk of complications:
 - **all children** who are immunocompromised, have a cochlear implant, craniofacial abnormalities (including cleft palate), developmental delay or Down syndrome, severe visual impairment, current hearing loss
 - **Aboriginal and Torres Strait Islander children** who reside in a remote community, first experience of otitis media at <6 months, family history of chronic suppurative otitis media (CSOM), current (or history of) perforation ⁽¹⁾.

Key points

- Acute otitis media (AOM) is predominantly a childhood infection, however, it may also occur in adults ⁽²⁾. The peak age prevalence is 6-18 months ⁽³⁾.
- AOM is commonly viral but may also be bacterial, or a combination of both ⁽⁴⁾.
- In the majority of paediatric cases, AOM is self-limiting and spontaneously resolves without the requirement for antibiotics ^(3, 4). The mainstay of treatment for AOM is conservative management with regular and adequate analgesia and aural hygiene. Antibiotics can be safely withheld in most cases and are often inappropriately prescribed ⁽⁴⁾.
- Pain alone is not sufficient to diagnose AOM; pneumatic otoscopy is required to diagnose and exclude other ear conditions with similar presentations⁽⁴⁾.
- Ear conditions are often very painful, particularly for young children, and it is important to approach with care ⁽¹⁾. If the patient will not allow for otoscopic examination due to pain, advice should be provided on appropriate analgesia and the patient referred to a medical practitioner.

When applying the information contained within this clinical practice guideline, pharmacists are advised to exercise professional discretion and judgement. The clinical practice guideline does not override the responsibility of the pharmacist to make decisions appropriate to the circumstances of the individual, in consultation with the patient and/or their carer.



Refer when

- The patient has 'red flag' warning signs
- A clear diagnosis of AOM cannot be made and when the TM cannot be visualised
- The patient is < 2 years of age

Treat (if clinically appropriate) and concurrently refer:

- The patient has otic discharge/otorrhea present
- Children presenting with signs and symptoms of a complication, or at a high risk of complications from AOM (refer to Table 1)
- The patient's symptoms do not improve and/or worsen after 48 hours of appropriate analgesia and supportive management
- Symptoms are recurrent (3 or more episodes in 6 months or 4 or more episodes in 12 months)
- The patient has persistent otitis media with effusion (glue ear).

Children who are prescribed antibiotic therapy must be concurrently referred to a medical practitioner for review.

Gather information and assess patient's needs

In both children and adults, the diagnosis of AOM is based on clinical history and ear examination using a pneumatic otoscope ^(1, 2, 4, 5).

- Children with AOM may present with ear pain and signs of systemic illness including fever, vomiting, diarrhoea, irritability, poor feeding and ear discharge within the preceding 6 weeks ⁽¹⁾. Tugging, holding or rubbing the ear may indicate ear pain in younger children ⁽⁴⁾.
- In children, a diagnosis of AOM requires the rapid onset (within 48 hours) of:
 - pain
 - signs of acute inflammation of the TM:
 - TM is bulging and opaque
 - TM may be red from inflammation or white from pus in the middle ear
 - signs of middle ear effusion
 - otorrhoea can be indicative of TM perforation and middle ear effusion if AOE has been excluded ^(1, 3, 4, 6)
 - otitis media with effusion is characterised by middle ear effusion without other signs and symptoms of AOM, where this is present longer than 3 months is suggestive of persistent otitis media with effusion (glue ear) ⁽⁴⁾
 - otorrhea that has been present for ≥ 2 weeks and a perforation is suggestive of chronic suppurative otitis media (CSOM) ⁽¹⁾.
- **A red TM or pain alone is not diagnostic of AOM** ⁽⁴⁾.
- In adults, unilateral symptoms are most common and include mild to severe ear pain and decreased hearing.
 - An upper respiratory tract infection (URTI) or allergic rhinitis will often precede the onset of AOM ⁽²⁾.
 - Other symptoms in adults include dizziness/vertigo (infrequent), systemic symptoms (particularly fever), partial or complete opacification of the TM, or erythema ⁽²⁾.
- In adults, the key signs of AOM are bulging and reduced mobility of the TM ⁽²⁾.

Patient history

Sufficient information should be obtained from the patient to assess the safety and appropriateness of any recommendations and medicines.

The patient history should consider:

- age
- whether the patient identifies as Aboriginal and Torres Strait Islander
- nature, severity and frequency of symptoms including pain, itch, discharge, hearing loss, a feeling of fullness and dizziness/vertigo
- onset and duration of symptoms including preceding illnesses

- recent history of URTI
- other presenting symptoms including temperature $> 38^{\circ}\text{C}$ or $< 35.5^{\circ}\text{C}$, rash, increased respiratory rate/distress, dehydration and/or reduced urine output, runny nose, sore throat or cough
- family history of ear conditions and hearing loss
- previous history of AOE, OM or other ear conditions:
 - hearing loss and recent hearing tests
 - patient age at first episode of ear conditions
 - under the care of an ENT specialist or audiologist
 - ear surgery, current or previous tympanostomy tubes (grommets)
- precipitating and relieving factors
- underlying medical conditions, e.g., immunocompromised (including diabetes), cochlear implant, craniofacial abnormalities, persistent hearing loss, that mean the patient is at high risk of experiencing AOM complications (refer to Table 1)
- delayed speech or language development (referral is required if this is evident and has not been assessed by a medical practitioner)
- social history including arrangements to support management at home, ability to access further care and to undertake appropriate self-care
- current medications (including prescribed medicines, vitamins, herbs, other supplements and over-the-counter medicines)
- frequency and method of ear cleaning
- medication and other strategies tried to treat current symptoms
- drug allergies/adverse drug effects
- immunisation status as per the Australian Immunisation Handbook.



Reminder

Pharmacists can access a range of clinical information in a patient's My Health Record, including details about current and past medication history, allergies and current medical conditions.

Assessment


- Conduct assessment of vital signs
- Examine for tender or enlarged cervical lymph nodes
- Conduct visual examination and palpation of mastoid area.

Ear examination using otoscopy

- Physical examination including otoscopy of both ears is required to differentiate between AOE, AOM and other ear conditions ^(2, 4, 5).

- Ear examination should be conducted in accordance with the [Queensland Health and Royal Flying Doctors Primary Clinical Care Manual](#). An ear differential diagnosis flowchart is also included ⁽¹⁾.
- Before performing the ear examination, the ear canal must be gently cleaned to remove any discharge with a tissue rolled up to a point (dry mopping) to visualise ear drum ^(1, 7). The [Remote Primary Health Care Manual – Clinical Procedure Manual](#) ⁽⁷⁾ and the [Queensland Health and Royal Flying Doctors Primary Clinical Care Manual](#) contain a step by step guide for dry mopping ⁽¹⁾.
- The TM must be visualised to confirm a diagnosis of AOM, although this can be difficult in the presence of extensive otorrhoea ⁽⁴⁾.
 - If the TM cannot be visualised in paediatric patients when they first present and antibiotics are not indicated, they should be advised on appropriate analgesia and either asked to return within 48 to 72 hours for reassessment, or to seek medical practitioner care if their condition deteriorates ⁽⁴⁾.
 - If the TM cannot be visualised in adult patients, they should be referred to a medical practitioner.
 - Perforation of the TM in children is not uncommon and may result in ear discharge and relief of pain ⁽¹⁾.
 - In AOM with perforation, the pharmacist should document the size and position of the perforation to allow ongoing assessment of progression of the condition and to guide use of therapies ⁽⁸⁾.
 - All medium to large perforations and perforations in the attic region of the TM (refer to Figure 1) must be referred to a medical practitioner.
- If pain levels allow, pneumatic otoscopy may be conducted to check for TM movement:
 - If the TM moves freely back and forward, perforation and fluid in the middle ear (AOM) can generally be excluded.
 - Reduced mobility of an intact TM is a good indication of the presence of middle ear fluid ⁽⁸⁾.
- Check for other symptoms if required (nose, throat, cough) ⁽¹⁾.

Figure 1. Guide for size of perforation of tympanic membrane ⁽⁸⁾



- TM “red flag” signs that may suggest a serious complication and require immediate referral to a medical practitioner include:
 - severe retraction of the TM
 - perforation or retraction in the attic region or near the edge of the TM

- crust, granulation or discharge in the attic region
- a dull white mass behind TM or chalky patches ⁽¹⁾.

Complications of AOM

Aboriginal and Torres Strait Islander children are generally at higher risk of complications arising from AOM. Complications in adults and non-First Nations children are less common ^(1, 6).

Children at high risk of complications are outlined in Table 1.

Table 1. Children at high risk of complications from AOM ⁽¹⁾

Table 1. Children at high risk of complications from AOM ⁽¹⁾	
Aboriginal and Torres Strait Islander children if: <ul style="list-style-type: none"> • residing in a remote community • < 2 years old • first episode of OM experience < 6 months of age • family history of CSOM • history (or current) perforation 	All children if: <ul style="list-style-type: none"> • immunocompromised • cochlear implant • craniofacial abnormalities including cleft palate • developmental delay, delayed speech, Down syndrome • severe visual impairment • current hearing loss

All children at high risk of complications must be referred to a medical practitioner for management. Antibiotic therapy may be provided concurrently to medical practitioner referral where clinically appropriate.

In rare cases, AOM infection may spread from the middle ear (intracranial spread) causing mastoiditis, petrositis or labyrinthitis ⁽⁶⁾.

Intracranial spread can cause meningitis or other intracranial complications that are slow to resolve, particularly in immunocompromised patients.

- Signs of intracranial complications include severe headache, confusion, focal neurologic signs, facial paralysis/palsy and dizziness/vertigo ⁽⁶⁾.

Management and treatment plan

Pharmacist management of patients with AOM involves:

- **supportive management:**
 - Aural hygiene and care at home in accordance with [The Royal Children's Hospital Melbourne Fact sheet: Ear infections and glue ear](#) and the [Australian Medicines Handbook: drugs for ear infections](#) ^(9, 10).
 - If discharge is present, advice and education regarding keeping ears dry and the use of dry mopping with tissue spears. The [Queensland Health and Royal Flying Doctors Primary Clinical Care Manual](#) contain a step by step guide for dry mopping ⁽¹⁾.

- **pharmacotherapy:**
 - Analgesia in accordance with the [Therapeutic Guidelines: Pharmacological management of acute pain](#).
 - Oral antibiotic therapy³ in accordance with the [Therapeutic Guidelines: Otitis media](#) ^(4, 8, 10).
- Consider immunisation where the patient has not received all routine vaccines in accordance with the [Australian Immunisation Handbook](#).

NB1: There is limited evidence to guide the management of AOM in adult patients. The Therapeutic Guidelines state that management of AOM is similar to children; adequate and regular analgesia is the primary treatment ⁽⁴⁾. As with paediatric patients, a shared decision-making approach is required to manage patient expectations of antibiotic treatment ⁽⁴⁾.

NB2: Paracetamol, Ibuprofen and Naproxen (in preparations containing 250mg or less) can be sold as Schedule 2 medicines.

NB3: Antibiotic therapy in paediatric patients should only be commenced in accordance with the Therapeutic Guidelines and the [Otitis Media Guidelines for Aboriginal and Torres Strait Islander People](#) for children with discharge/otorrhoea (from a small perforation), children with symptoms that have not improved or worsened after 48 hours and Aboriginal and Torres Strait Islander children ^(1, 4, 8). Children who are prescribed antibiotic therapy must be concurrently referred to a medical practitioner for review.

Confirm management is appropriate

Pharmacists must consult the Therapeutic Guidelines, Australian Medicines Handbook and other relevant references including the Otitis Media Guidelines for Aboriginal and Torres Strait Islander People (if applicable) to confirm that the treatment recommendation is appropriate, including for:

- contraindications and precautions
- drug interactions
- pregnancy and lactation.

Communicate agreed management plan

Comprehensive advice and counselling (including supporting written information when required) as per the Australian Medicines Handbook and other relevant references should be provided to the patient regarding:

- individual product and medicine use (e.g., dosing)
- how to manage adverse effects
- when to seek further care and/or treatment (including recognising patient deterioration)
- when to return to the pharmacist for clinical review.

It is the pharmacist's responsibility to ensure the suitability and accuracy of any resources and information provided to patients (and parents/caregivers if applicable) and to ensure compliance with all copyright conditions.

The agreed management plan should be shared with members of the patient's multidisciplinary healthcare team, with the patient's consent.

Patient resources/ information

- [The Royal Children's Hospital Melbourne Fact sheet: Ear infections and glue ear](#) ⁽⁹⁾.

General advice

AOM symptoms should start to improve within 48 to 72 hours after onset, regardless of whether antibiotic treatment is required, but full resolution of symptoms may take 8 days or longer for effusion ⁽⁴⁾.

- Middle ear effusion that persists for several weeks after AOM is common and is not a sign of treatment failure ⁽¹⁰⁾
- An acute perforation and discharge should resolve within 10-14 days ⁽¹⁾.

Patients commenced on antibiotics who have been concurrently referred to a medical practitioner should be advised (or their parents/caregivers if applicable) to seek further urgent care where:

- the response has been inadequate after 48-72 hours of commencing antibiotic treatment
- the patient's condition deteriorates, including if they develop systemic signs and symptoms or if their symptoms worsen
- otorrhea continues for ≥ 2 weeks ⁽⁴⁾.

For all patients, it is important to avoid local trauma to ear canal, including not using cotton ear buds and dry mopping with a tissue spear instead ⁽¹⁾. The [Remote Primary Health Care Manual – Clinical Procedure Manual](#) ⁽⁷⁾ and the [Queensland Health and Royal Flying Doctors Primary Clinical Care Manual](#) contain a step by step guide for dry mopping ⁽¹⁾.

Pharmacists should also provide advice on the importance of preventing ear infections in children to avoid associated hearing loss, poor language and social skills ⁽⁸⁾. Preventative advice that can be provided by pharmacists includes:

- ensuring vaccinations against *S. pneumoniae*, *H. influenzae* type B, and influenza are up to date
- minimising exposure to smoke e.g., cigarettes and woodfires
- encourage personal hygiene e.g., frequent hand and face washing and drying, nose blowing techniques ^(4, 8, 10, 11).

Decongestants, antihistamines and corticosteroids are not effective and should not be recommended for adult or paediatric patients ^(3, 10).

Antibiotics have limited benefits for most paediatric patients and are not indicated for most children presenting with mild infection ⁽⁴⁾.

- The use of antibiotics in paediatric patients almost doubles the incidence of diarrhoea, vomiting and rash ⁽¹⁰⁾
- Pharmacists should address misconceptions regarding antibiotic effectiveness and discuss the relative harms and benefits of treatment ⁽³⁾.

Clinical review

Clinical review should occur in line with recommendations in the Therapeutic Guidelines, the Queensland Health and Royal Flying Doctors Primary Clinical Care Manual, and other relevant guidelines.

- Patients with an intact TM are recommended to be reviewed at 3 months to check for persistent fluid behind the TM; if fluid is still present or the patient experiences a recurrence of AOM during this time, they must be referred to a medical practitioner ⁽¹⁾.
- Patients with small acute perforations are recommended to be reviewed between **4-6 weeks** to assess healing. If the perforation is still evident, the patient must be referred to a medical practitioner ⁽¹⁾.

The decision regarding the timing and necessity of a clinical review should be determined on an individual basis, considering the patient's (or parents/caregivers) ability and available support to manage the condition effectively. The pharmacist should discuss the importance of clinical review with the patient and/or their caregiver, including the timing of review, and where review by a medical practitioner is indicated.

For children under the age of 5, the improvement of the condition and the presence of complications (e.g., parent/carer concerns related to the child's speech and language) should be closely monitored ⁽¹²⁾.

For patients who have already been referred to a medical practitioner following initial assessment and/or management by the pharmacist, further clinical review by the pharmacist is generally not required.



Pharmacist resources

- Therapeutic Guidelines: Antibiotic
 - Otitis media
- Therapeutic Guidelines: Pain and analgesia
 - Pharmacological management of acute pain
- Australian Medicines Handbook
 - Otitis Media
 - Drug for ear infections
 - NSAIDs
- Australian Immunisation Handbook:
 - [Pneumococcal disease](#)
 - [Influenza \(flu\)](#)
 - [Haemophilus influenzae type b \(Hib\)](#)
- MSD Manual (Professional version) - [Otitis media](#)
- Remote Primary Health Care Manual:
 - [Clinical Procedure Manual](#)
 - [CARPA Standard Treatment Manual](#)
- Three minute tool kit (UK) - [Spotting the Sick Child](#)
- Queensland Health and Royal Flying Doctors Service (Queensland branch) [Primary Clinical Care Manual 11th edition 2022](#)
- The Royal Children's Hospital - [Clinical Practice Guidelines – acute otitis media](#)
- [2020 Otitis Media Guidelines for Aboriginal and Torres Strait Islander Children](#)
- McGovern Medical School - [Ear disease photo book](#)

Document version number	Date	Comments
Version 1.0	01.02.2024	
Version 1.1	11.11.2024	Administrative update

Copyright

© The State of Queensland 2024

The Queensland Government supports and encourages the distribution of its material.

Unless otherwise noted, all copyright material in this publication is licensed under a [Creative Commons Attribution 4.0 International licence \(CC BY 4.0\)](#).



Material in this publication may be reproduced in whole or in part for study or training purposes subject to the inclusion of an acknowledgement of the source. It may not be reproduced for commercial usage or sale. Reproduction for purposes other than those indicated above requires written permission from Queensland Health. © Queensland Health 2024.

You are free to use copyright material available in this publication that is covered by a CC BY licence in line with the licence terms. You must keep the copyright notice on the copyright material and attribute the State of Queensland as the source of the copyright material.

Note: Creative Commons licences relate to copyright material, not to other forms of intellectual property. Some material accessible in this publication may not be available under a Creative Commons licence.

Disclaimer

As far as practicable, material for which the copyright is owned by a third party will be clearly labelled. The Queensland Government has made all reasonable efforts to ensure that this material has been reproduced with the consent of the copyright owners.

The materials available in this publication are distributed by the Queensland Government as an information source only.

To the maximum extent permitted by law, the State of Queensland makes no statement, representation, or warranty about the quality, accuracy, context, completeness, availability or suitability for any purpose of, and you should not rely on, any materials available in this publication.

Despite best efforts, the State of Queensland makes no warranties that the materials available on or through any links provided in this publication are free of infection by computer viruses or other contamination, to the maximum extent permitted by law.

The Queensland Government disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you or any other person might incur for any reason including as a result of the materials available in this publication being in any way inaccurate, out of context, incomplete, unavailable, not up to date or unsuitable for any purpose.

References

1. Queensland Health and Royal Flying Doctor Service (Queensland Section). Primary Clinical Care Manual. Cairns: Office of Rural and Remote Health, Queensland Government; 2022 [cited 2022 September 18]. Available from: <https://www.health.qld.gov.au/rrcsu/clinical-manuals/primary-clinical-care-manual-pccm>.
2. Limb C, Lustig L, Durand M. Acute otitis media in adults: Wolters Kluwer; 2022 [cited 2022 April 12]. Available from: <https://www.uptodate.com/contents/acute-otitis-media-in-adults>.
3. The Royal Children's Hospital. Clinical practice guidelines: acute otitis media Melbourne: RCHM; 2021 [cited 2022 April 14]. Available from: https://www.rch.org.au/clinicalguide/guideline_index/Acute_otitis_media/.
4. Therapeutic Guidelines: Antibiotic (Otitis media). Melbourne: Therapeutic Guidelines Limited; 2021 [cited 2022 April 12]. Available from: <https://tgldcdp.tg.org.au/viewTopic?topicfile=otitis-media>.
5. Therapeutic Guidelines: Antibiotic (Otitis externa). Melbourne: Therapeutic Guidelines Limited; 2021 [cited 2022 April 14]. Available from: <https://tgldcdp.tg.org.au/viewTopic?topicfile=otitis-externa>.
6. Miyamoto R. MSD Manual Professional Version: Otitis Media (Acute): Merck & Co; 2022 [cited 2022 June 24]. Available from: <https://www.msdmanuals.com/en-au/professional/ear,-nose,-and-throat-disorders/middle-ear-and-tympanic-membrane-disorders/otitis-media-acute?query=otitis%20media>.
7. Remote Primary Health Care Manuals. Clinical Procedures Manual. Alice Springs: Centre for Remote Health; 2017 [cited 2022 June 23]. Available from: <https://remotephcmmanuals.com.au/>.
8. Leach AJ MP, Coates HLC, et al.,. Otitis media guidelines for Australian Aboriginal and Torres Strait Islander children. Northern Territory: Menzies School of Health Research; 2020.
9. The Royal Children's Hospital General Medicine Emergency and Otolaryngology departments. Ear infections and glue ear Melbourne: RCHM; 2018 [cited 2022 March 18]. Available from: https://www.rch.org.au/kidsinfo/fact_sheets/Ear_infections_and_otitis_media/.
10. Australian Medicines Handbook: Drugs for ear infections. Adelaide: Australian Medicines Handbook Pty Ltd; 2022 [cited 2022 September 18]. Available from: <https://amhonline.amh.net.au/chapters/ear-nose-throat-drugs/drugs-ear-infections?menu=vertical>.
11. Remote Primary Health Care Manuals. CARPA Standard Treatment Manual. Alice Springs: Centre for Remote Health; 2017 [cited 2022 June 23]. Available from: <https://remotephcmmanuals.com.au/>.
12. Queensland Health; Royal Flying Doctors Service (Queensland Section) and Apunipima Cape York Health Council. Chronic Conditions Manual: Prevention and Management of Chronic Conditions in Rural and Remote Australia. Cairns: The Rural and Remote Clinical Support Unit, Torres and Cape Hospital and Health Service; 2020.