



Iodinated Contrast Injection- Patients with Renal Impairment

Facility: _____

(Affix identification label here)

URN: _____

Family name: _____

Given name(s): _____

Address: _____

Date of birth: _____

Sex: M F I

A. Interpreter / cultural needs

An Interpreter Service is required? Yes No

If Yes, is a qualified Interpreter present? Yes No

A Cultural Support Person is required? Yes No

If Yes, is a Cultural Support Person present? Yes No

B. Procedure

The following will be performed (*Doctor/doctor delegate to document – include site and/or side where relevant to the procedure*)

.....
.....
.....

Iodinated 'Contrast' is colourless liquid containing iodine. It is injected into your blood stream to allow your organs to be seen more clearly on x-ray.

Contrast is not a dye. It does not stain the inside of your body.

You have been identified as having renal impairment (weakened kidneys) through the questions asked by staff and by having a blood test.

Giving the Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).

C. Risks of Iodinated Contrast

In recommending the administration of Iodinated Contrast for this procedure, the doctor believes the benefits to you from having Contrast exceed the risks involved.

The risks and complications with Contrast can include but are not limited to the following:

Common risks and complications include:

- No known common risks.

Less common risks and complications include:

- Injected Contrast may leak outside of the blood vessel, under the skin and into the tissues. This may require treatment. In very rare cases, further surgery could be required if the skin breaks down.
- Acute Renal Failure occurs when one or both of your kidneys suddenly stop working. Failure can last for days or weeks. It may take the kidneys a long time to regain their previous level of function and you may require dialysis to filter your blood during this time. There is a risk that kidneys could be permanently damaged. To reduce this risk the smallest possible dose of Contrast will be given.

- The injection may not be possible due to medical and/or technical reasons.

Rare risks and complications include:

- Allergic reactions occur within the first hour with most happening in the first 5 minutes. Late reactions have been known to occur up to a week after the injection.

Note: Allergy to topical iodine and/or seafood does not imply an allergy to Iodinated Contrast
The reactions vary from:

Mild – hives, sweating, sneezing, coughing, nausea.

Moderate – wide spread hives, headaches, facial swelling, vomiting, shortness of breath.

Severe – Severe reactions are rare but include: life-threatening heart palpitations, very low blood pressure, throat swelling, fits and/or cardiac arrest.

- Death as a result of Iodinated Contrast is very rare.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....



(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

Facility:

Iodinated Contrast Injection- Patients with Renal Impairment

D. Patient consent

I acknowledge that the doctor/doctor delegate has explained the proposed procedure.

I understand;

- the risks and complications, including the risks that are specific to me.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor/doctor delegate or my Acute Resuscitation Plan.
- a doctor/doctor delegate undergoing further training may conduct this procedure.

I have been given the following Patient Information Sheet/s:

Iodinated Contrast Injection - Patients with Renal Impairment

- I was able to ask questions and raise concerns with the doctor/doctor delegate about the proposed procedure and its risks. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time including after I have signed this form but, preferably following a discussion with my doctor/doctor delegate.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.
- I understand that Queensland Health may release my relevant de-identified information obtained from this and related procedures for education and training of health professionals.

On the basis of the above statements,

I request to have the injection

Name of Patient:

Signature:

Date:

Patients who lack capacity to provide consent

Consent must be obtained from a substitute decision maker/s in the order below.

Does the patient have an Advance Health Directive (AHD)?

Yes ▶ Location of the original or certified copy of the AHD:

No ▶ Name of Substitute Decision Maker/s:

Signature:

Relationship to patient:

Date: PH No:

Source of decision making authority (tick one):

- Tribunal-appointed Guardian
- Attorney/s for health matters under Enduring Power of Attorney or AHD
- Statutory Health Attorney
- If none of these, the Adult Guardian has provided consent. Ph 1300 QLD OAG (753 624)

E. Doctor/delegate Statement

I have explained to the patient all the above points under the Patient Consent section (D) and I am of the opinion that the patient/substitute decision-maker has understood the information.

Name of Doctor/delegate:

Designation:

Signature: Date

F. Interpreter's statement

I have given a sight translation in

.....

(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

Name of Interpreter:

Signature: Date

DO NOT WRITE IN THIS BINDING MARGIN

1. What is Iodinated Contrast?

The medical imaging procedure your doctor has asked you to have uses Iodinated 'Contrast' (once called X-ray dye). Contrast is a colourless liquid which includes iodine. Contrast is injected into your blood stream to allow your organs to be seen more clearly on x-rays. Contrast is not a dye. It does not stain the inside of your body. Your doctor needs to use Contrast to be able to get all the information needed to assist with your diagnosis.

This information sheet must be read together with the information sheet of the procedure you are booked for (if you do not have this information sheet please ask for one).

2. During the procedure

When the Contrast is injected you may feel:

- A very warm or 'flushed' feeling over your body, this may also make you think you have passed urine. You will *not* pass urine – it is only a feeling.
- A 'metallic' taste or smell may also happen. This usually lasts less than a minute.

3. After the procedure

It is recommended that you drink 2 to 4 glasses of water after your procedure to help flush the Contrast from your body.

Contrast does not affect your ability to carry out normal activities; you should be able to continue with your day as normal.

4. Precautions

Contrast is not suitable for some people; you will be asked a series of questions before it is given to you. Your answers allow staff to identify any risk factors that you may have.

- Please tell the staff if you are or suspect you might be pregnant or are breastfeeding.

Kidney function

- Contrast is removed from your blood by your kidneys through your urine. It is easily removed from the body of people who have normal kidney function.
- Giving Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).
- You may be asked to have a simple blood test to find out the level of their kidney function.

Diabetic Drug interactions - Metformin

(Other Drug names: Avandamet, Diabex, Diaformin, Formet, Glucohexal, Glucomet, Glucophage, Glucovance, Metforbell)

If kidneys suffer damage from the Contrast then the kidneys may not be able to remove Metformin from the body. It is safer to briefly stop taking Metformin when having Contrast. Staff will inform you when to stop and when it is safe to take Metformin again.

Contact your GP to monitor your diabetes if you are told to stop your Metformin.

5. What are the risks of Iodinated Contrast?

The risks and complications with this injection can include but are not limited to the following.

Common risks and complications include:

- No known common risks.

Less common risks and complications include:

- Injected Contrast may leak outside of the blood vessel, under the skin and into the tissue. This may require treatment. In very rare cases, further surgery could be required if the skin breaks down.
- Acute Renal Failure occurs when one or both of your kidneys suddenly stop working. Failure can last for days or weeks. It may take the kidneys a long time to regain their previous level of function and you may require dialysis to filter your blood during this time. There is a risk your kidneys could be permanently damaged. To reduce this risk the smallest possible dose of Contrast will be given.
- The injection may not be possible due to medical and/or technical reasons.

Rare risks and complications include:

- Allergic reactions occur within the first hour with most happening in the first 5 minutes. Late reactions have been known to occur up to a week after the injection.

Note: Allergy to topical iodine and/or seafood does not imply an allergy to Iodinated Contrast.

The reactions vary from:

Mild – hives, sweating, sneezing, coughing, nausea.

Moderate – wide spread hives, headaches, facial swelling, vomiting, shortness of breath.

Severe – Severe reactions are rare but include: life-threatening heart palpitations, very low blood pressure, throat swelling, fits and/or cardiac arrest.

- Death as a result of Iodinated Contrast is *very* rare.

6. What are the safety issues when you leave the hospital?

Go to your nearest Emergency Department or GP if you become unwell.