

Queensland Community Pharmacy Scope of Practice Pilot

Acute Nausea and Vomiting - Clinical Practice Guideline

Guideline Overview





'Red flag' warning signs at patient presentation that necessitate referral to a medical practitioner:

- Prolonged vomiting (longer than 24-48 hours)
- Patient looks very unwell/very drowsy
- Significant recent weight loss
- Abdominal distension or tenderness
- Rectal bleeding
- Green, bile or blood/'coffee grounds' vomit
- Fever, neck stiffness, confusion
- Severe headache, altered or loss of consciousness
- Isolated vomiting, lack of nausea (nausea without vomiting is not uncommon)
- History of head injury/trauma
- Increased blood glucose levels.

Key points

- It is essential to consider the clinical context when assessing patients with nausea and vomiting, including the possibility of life-threatening causes, before developing an individualised management plan ^(1, 2).
- Causes of nausea and vomiting can stem from the gastrointestinal (GI) tract, the central nervous system (CNS), or result from a systemic condition, and can often be attributed to an identifiable cause, most commonly acute gastroenteritis, food poisoning or other toxins, migraine, cannabis use, motion sickness or pregnancy ^(1, 3).
- An oral antiemetic may be considered for symptom relief, although spontaneous improvement of acute nausea and vomiting is likely ⁽¹⁾.

When applying the information contained within this clinical practice guideline, pharmacists are advised to exercise professional discretion and judgement. The clinical practice guideline does not override the responsibility of the pharmacist to make decisions appropriate to the circumstances of the individual, in consultation with the patient and/or their carer.



Refer when

- The patient has 'red flag' warning signs
- The patient is aged under 18 years
- The patient has diabetes or is immunocompromised
- The patient has recently returned from overseas
- The nausea and vomiting are post-operative, radiation-induced or suspected to be drug-induced
- Intramuscular administration of an antiemetic is indicated
- The nausea and vomiting is chronic (defined by 4 weeks or more of symptoms) ⁽⁴⁾
- The patient has recurring episode of acute nausea and vomiting
- The symptoms have not resolved within 24-48 hours of first commencing
- The patient's condition and/or symptoms worsen.

Treat (if clinically appropriate) and concurrently refer:

- The patient is pregnant

Gather information and assess patient's needs

The assessment approach focuses on identifying the cause (or excluding significant underlying diseases) with a view to expectant management or directing specific treatment.

Patient history

Sufficient information should be obtained from the patient to identify the likely cause of the nausea/vomiting, and assess the safety and appropriateness of any recommendations and medicines.

The patient history should consider:

- age
- pregnancy and lactation status (if applicable)
- nature, severity and frequency of symptoms
- food and/or drink consumed in the last 24 hours
- onset and duration of symptoms, including timing in relation to eating/drinking
- recent head trauma or injury
- similar symptoms in close contacts
- exposure to toxins, poisons, bites or stings
- recent travel
- other symptoms e.g., chest pain or heartburn, headache, dizziness, fever, abdominal pain, last bowel motion, diarrhoea, constipation, dysuria, or frequency of urination
- precipitating and relieving factors
- underlying medical conditions e.g., diabetes or recent surgery
- current medications (including prescribed medicines, vitamins, herbs, other supplements, and over-the-counter medicines), including initiation of new medicines

- medication tried to treat current symptoms
- alcohol and drug history, including recent illicit drug or alcohol use.



Reminder

Pharmacists can access a range of clinical information in a patient's My Health Record, including details about current and past medication history, allergies and current medical conditions.

Examination

- Where appropriate, conduct assessment of vital signs.
- Assess hydration status as per the [Therapeutic Guidelines: Gastrointestinal – Assessment of acute gastroenteritis](#) ⁽⁵⁾.
- Examine for abdominal tenderness or distention, particularly areas where hernias commonly present excluding the groin.

Consider possible causes of nausea and vomiting

A non-exhaustive list of possible causes of acute nausea and vomiting are presented in Appendix 1. When in doubt of the cause of the nausea and vomiting, the pharmacist should refer the patient to a medical practitioner.

Management and treatment plan

Before starting treatment, it is important to identify, treat or remove the cause of the nausea and vomiting, and ensure adequate hydration ⁽⁶⁾.

Pharmacist management of acute nausea and vomiting includes:

- **supportive management for gastroenteritis:**
 - Including oral rehydration therapy to prevent or correct dehydration, in accordance with the [Therapeutic Guidelines: Acute gastroenteritis](#) ⁽⁵⁾.
- **pharmacotherapy:**
 - In accordance with the [Therapeutic Guidelines: Nausea and vomiting](#), excluding ondansetron, ^{1,2,3 (1)}, limited to 24 hours supply of oral regimens for schedule 4 medicines.

NB1: Prochlorperazine is available as a Schedule 3 medicine limited to a pack size of 10 tablets (5 mg). If there is a need for further medication, the patient should be referred to a medical practitioner.

NB2: Promethazine is available as a Schedule 3 in a pack size of 25 or 50 tablets (25 mg).

NB3: Ondansetron is indicated for the prevention and treatment of nausea and vomiting induced by cytotoxic therapy and radiotherapy, and the prevention of post-operative

nausea and vomiting. Its use for any other indication is 'off-label' and not included in the Pilot.

Confirm management is appropriate

Pharmacists must consult the Therapeutic Guidelines, Australian Medicines Handbook and other relevant references to confirm that the treatment recommendation is appropriate, including for:

- contraindications and precautions
- drug interactions
- pregnancy and lactation.

Communicate agreed management plan

Comprehensive advice and counselling (including supporting written information when required) as per the Australian Medicines Handbook and other relevant references, should be provided to the patient regarding:

- individual product and medicine use e.g., dosing and frequency
- supportive care, including the importance of rehydration
- how to manage adverse effects
- the impact of vomiting on the absorption and effectiveness of other regular medications
- when to seek further care and/or treatment, including recognising 'red flags' and dehydration.

It is the pharmacist's responsibility to ensure the suitability and accuracy of any resources and information provided to patients (and parents/caregivers if applicable) and to ensure compliance with all copyright conditions.

The agreed management plan should be shared with members of the patient's multidisciplinary healthcare team, with the patient's consent.

Clinical review

Clinical review with the pharmacist is generally not required. If the condition does not improve or resolve within 24-48 hours, the patient should be advised to see a medical practitioner for further investigation.



Pharmacist resources

- Therapeutic Guidelines:
 - Nausea and vomiting
 - Acute gastroenteritis
- Australian Medicines Handbook - Antiemetics
- MSD Manual (Professional version):
 - [Nausea and vomiting](#)
 - [Acute abdominal pain](#)
- SOMANZ - [Guideline for the Management of Nausea and Vomiting in Pregnancy and Hyperemesis Gravidarum](#)
- Queensland Health and Royal Flying Doctors Service (Queensland branch) - [Primary Clinical Care Manual 11th edition 2022](#)
- Australian Family Physician (RACGP) - [Nausea and vomiting in adults: A diagnostic approach.](#)
- Food safety:
 - [Outbreak management](#)
 - [Make a food safety complaint online](#)

Appendix 1 - Possible causes of nausea and vomiting

| Causes of nausea and vomiting ^(1, 3, 7, 8) | | |
|---|--|---|
| Condition | Presentation and suggestive findings | Action |
| Gastrointestinal disorders | | |
| Adynamic ileus | <ul style="list-style-type: none"> Abdominal distention Increased risk after surgery (up to 72 hours post-surgery) Severe illness or electrolyte abnormality | Referral to a medical practitioner if suspected. |
| Bowel obstruction | <ul style="list-style-type: none"> Abdominal distention, severe constipation and tympany to percussion Vomiting is often bilious History of abdominal surgery (visible surgical scars) or hernia | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| Gastroenteritis – viral or bacterial/ food poisoning | <ul style="list-style-type: none"> Vomiting accompanied by diarrhoea Normal abdomen when examined Food poisoning apparent based on history | Self-limiting – May be managed under this guideline. For suspected food poisoning, consider notification to local public health unit or make a food safety complaint online . |
| Gastroparesis | <ul style="list-style-type: none"> Vomiting of partially digested food within a few hours of consumption Often occurs in people with diabetes or after abdominal surgery | Referral to a medical practitioner is required if suspected. |
| Hepatitis | <ul style="list-style-type: none"> Mild to moderate nausea (sometimes vomiting) over many days Jaundice Malaise Slight liver tenderness is sometimes present | Referral to a medical practitioner is required if suspected. |
| Peritonitis or other acute abdominal condition e.g., pancreatitis, appendicitis | <ul style="list-style-type: none"> Severe abdominal pain that may be vague and nauseating (visceral), sharp and localised (somatic) or pain perceived distant to its source (referred) Other warning signs (see Page 1) may be present | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |

| Causes of nausea and vomiting ^(1, 3, 7, 8) | | |
|--|--|---|
| Toxic ingestion including alcohol | <ul style="list-style-type: none"> Generally apparent based on history | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| Central nervous system disorders | | |
| Cannabis use (cannabis hyperemesis syndrome) | <ul style="list-style-type: none"> Ongoing nausea, vomiting and indigestion that occurs with chronic cannabis use | Symptoms resolve after stopping cannabis use and may be relieved with a hot shower. Referral to a medical practitioner and other services if required |
| Closed head injury or concussion | <ul style="list-style-type: none"> Generally apparent based on history | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| CNS haemorrhage | <ul style="list-style-type: none"> Sudden onset of headache Change in mental status Meningeal signs including photophobia, neck stiffness and seizures may be present | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| CNS infection e.g., meningitis | <ul style="list-style-type: none"> Gradual onset of headache Change in mental status Meningeal signs including photophobia, neck stiffness and seizures may be present Petechial rash may be present | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| Increased intracranial pressure | <ul style="list-style-type: none"> Headache Change in mental status Focal neurological deficits sometimes present | Immediate referral to a medical practitioner (including emergency service) is required if suspected. |
| Labyrinthitis | <ul style="list-style-type: none"> Vertigo, nystagmus with symptoms worsened by motion Tinnitus sometimes present | Referral to a medical practitioner is required if suspected. |
| Migraine | <ul style="list-style-type: none"> Headache, with or without aura or photophobia Patient will usually have a history | Standard pharmacy treatment and referral |

| Causes of nausea and vomiting ^(1, 3, 7, 8) | | |
|--|--|--|
| | | to a medical practitioner. |
| Motion sickness | <ul style="list-style-type: none"> Generally apparent based on history | Standard pharmacy treatment. |
| Systemic conditions | | |
| Advanced cancer, chemotherapy or radiation exposure | <ul style="list-style-type: none"> Apparent based on history | Referral to a medical practitioner is required. |
| Diabetic ketoacidosis (DKA) | <ul style="list-style-type: none"> History of diabetes, although may occur without a known history of diabetes Polyuria, polydipsia, hyperventilation, abdominal pain and/or impaired consciousness Significant dehydration often present Patients with diabetes should be advised to check their blood glucose levels and test ketones in blood or urine ⁽⁹⁾ | Immediate referral to a medical practitioner (including an emergency service) is required if this is suspected, including if ketones are 0.6 or higher. |
| Drug adverse effect or toxicity | <ul style="list-style-type: none"> Generally apparent based on history Assess severity and probable cause | <p>If nausea and vomiting is a known adverse effect, referral to the original prescriber is required.</p> <p>If suspected overdose or serious reaction, immediate referral to a medical practitioner (including emergency services) is required.</p> |
| Liver or renal failure | <ul style="list-style-type: none"> Generally apparent based on history Asterixis Jaundice in advanced liver disease often present Uremic odour in renal failure often present | Immediate referral to a medical practitioner (including emergency services) is required if suspected. |
| Severe pain | <ul style="list-style-type: none"> Multiple causes and presentations | Immediate referral to a medical practitioner (including emergency services) is required. |

| Causes of nausea and vomiting ^(1, 3, 7, 8) | | |
|--|---|--|
| Addisonian (acute adrenal) crisis | <ul style="list-style-type: none"> History of adrenal insufficiency, trauma, acute stress, illness or recent surgery, or abrupt cessation of glucocorticoid therapy Symptoms of acute circulatory failure (hypotension, confusion) Other symptoms may include weakness, loss of consciousness, dizziness, fever, sudden onset of pain in lower back or legs, high heart rate | Immediate referral to a medical practitioner (including emergency services) is required. |
| Other causes | | |
| Pregnancy | <ul style="list-style-type: none"> Can occur at any time of the day or night. May be triggered by food or odours Patient may not be aware they are pregnant | Referral to a medical practitioner is required and non-pharmacological therapies may be recommended as standard pharmacist care. |
| Urinary Tract Infection | <ul style="list-style-type: none"> Generally apparent based on history Presence of dysuria, urinary urgency and frequency Suprapubic tenderness or discomfort | Referral to a medical practitioner or an appropriate pharmacy service is required. |

| Document version number | Date | Comments |
|--------------------------------|-------------|-----------------------|
| Version 1.0 | 01.02.2024 | |
| Version 1.1 | 11.11.2024 | Administrative update |

Copyright

© The State of Queensland 2024

The Queensland Government supports and encourages the distribution of its material.

Unless otherwise noted, all copyright material in this publication is licensed under a [Creative Commons Attribution 4.0 International licence \(CC BY 4.0\)](#).



Material in this publication may be reproduced in whole or in part for study or training purposes subject to the inclusion of an acknowledgement of the source. It may not be reproduced for commercial usage or sale. Reproduction for purposes other than those indicated above requires written permission from Queensland Health. © Queensland Health 2024.

You are free to use copyright material available in this publication that is covered by a CC BY licence in line with the licence terms. You must keep the copyright notice on the copyright material and attribute the State of Queensland as the source of the copyright material.

Note: Creative Commons licences relate to copyright material, not to other forms of intellectual property. Some material accessible in this publication may not be available under a Creative Commons licence.

Disclaimer

As far as practicable, material for which the copyright is owned by a third party will be clearly labelled. The Queensland Government has made all reasonable efforts to ensure that this material has been reproduced with the consent of the copyright owners.

The materials available in this publication are distributed by the Queensland Government as an information source only.

To the maximum extent permitted by law, the State of Queensland makes no statement, representation, or warranty about the quality, accuracy, context, completeness, availability or suitability for any purpose of, and you should not rely on, any materials available in this publication.

Despite best efforts, the State of Queensland makes no warranties that the materials available on or through any links provided in this publication are free of infection by computer viruses or other contamination, to the maximum extent permitted by law.

The Queensland Government disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs you or any other person might incur for any reason including as a result of the materials available in this publication being in any way inaccurate, out of context, incomplete, unavailable, not up to date or unsuitable for any purpose.

References

1. Therapeutic Guidelines: Gastrointestinal (Nausea and vomiting). Melbourne: Therapeutic Guidelines Limited; 2021 [cited 2022 September 23]. Available from: https://tgldcdp.tg.org.au/viewTopic?topicfile=c_GIG_Assessment-and-causes-of-nausea-and-vomitingtopic_1.
2. Queensland Health and Royal Flying Doctor Service (Queensland Section). Primary Clinical Care Manual. Cairns: Office of Rural and Remote Health, Queensland Government; 2022 [cited 2022 September 23]. Available from: https://www.health.qld.gov.au/_data/assets/pdf_file/0011/1160003/PCCM_11th_ed.pdf.
3. Gotfried J. MSD Manual Professional Version: Nausea and Vomiting; Merck & Co; 2022 [cited 2022 June 8]. Available from: <https://www.msdmanuals.com/professional/gastrointestinal-disorders/symptoms-of-gastrointestinal-disorders/nausea-and-vomiting?query=nausea%20and%20vomiting>.
4. Lacy BE, Parkman HP, Camilleri M. Chronic nausea and vomiting: evaluation and treatment. *Am J Gastroenterol*. 2018;113(5):647-59.
5. Therapeutic Guidelines: Gastrointestinal (Acute gastroenteritis). Melbourne: Therapeutic Guidelines Limited; 2021 [cited 2022 September 23]. Available from: https://tgldcdp.tg.org.au/viewTopic?etgAccess=true&guidelinePage=Gastrointestinal&topicfile=c_GIG_Assessment-of-acute-gastroenteritistopic_1&guidelinename=Gastrointestinal§ionId=c_GIG_Assessment-of-acute-gastroenteritistopic_1#c_GIG_Assessment-of-acute-gastroenteritistopic_1.
6. Australian Medicines Handbook: Antiemetics. Adelaide: Australian Medicines Handbook Pty Ltd; 2022 [cited 2022 September 23]. Available from: <https://amhonline.amh.net.au/chapters/gastrointestinal-drugs/antiemetics?menu=vertical>.
7. Metz A, Hebbard G. Nausea and Vomiting in Adults: A diagnostic approach. *Australian Family Physician*. 2007;36(9).
8. Ansari P. MSD Manual Professional Version: Acute Abdominal Pain; Merck & Co; 2021 [cited 2022 June 16]. Available from: <https://www.msdmanuals.com/professional/gastrointestinal-disorders/acute-abdomen-and-surgical-gastroenterology/acute-abdominal-pain>.
9. Statewide Diabetes Network Queensland Health. Management of diabetic ketoacidosis in adults. Brisbane: The State of Queensland; 2015 [cited 2022 September 9]. Available from: <https://clinicalexcellence.qld.gov.au/resources/diabetes-resources/diabetic-ketoacidosis>.