Queensland	(Affix identification	on label here)
Government	URN:	
	Family name:	
Fractured Forearm	Given name(s):	
	Address:	
Facility:	Date of birth:	Sex: M F I
A. Interpreter / cultural needs	Specific risks:	

1 domey.	
A. Interpreter / cultural needs	
An Interpreter Service is required?	No
If Yes, is a qualified Interpreter present?	No
A Cultural Support Person is required?	No
If Yes, is a Cultural Support Person present? Yes	No
B. Condition and treatment	
The doctor has explained that you have the following condition: (Doctor to document in patient's own wo	_
This condition requires the following procedure. (Doctor to document - include site and/or side whe relevant to the procedure)	re
Left arm	

The following will be performed:

Right arm

This procedure is the fixation of the fractured forearm with the placement of plate and screws or nails.

☐ Yes ☐ No

C. Risks of a fractured forearm

There are risks and complications with this procedure. They include but are not limited to the following. General risks:

- Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

- Numbness associated with the use of tourniquet with nerve and muscle damage at the site where the tourniquet was placed. This may be temporary or permanent.
- Skin death under the tourniquet, which may require further dressings and / or surgery and skin grafting.
- Damage to nerves and/or vessels resulting in numbness and/or weakness. This may temporary or permanent. Another operation may be required to correct the situation.
- Stiffness in the joints may occur above and below the break in the arm.
- Build up of pressure in the muscle compartment, which can require re-operation.
- Inability to close the edges of the wound together.
 This may require skin grafts.
- The bones do not knit together properly. This may require further surgery and/or bone graft.
- The bones heal with a bend in the bones and/or alignment.
- The implants may need to be removed in future months or years. This will require further surgery.
- Abnormal pain response to surgery with worsening of pain and disability.
- The surgical cut may cause changes to the sensation and colour of the limb.
- In some people, healing of the wound may be abnormal and the wound can be thickened and red and the scar may be painful.

D.	Significant risks and procedu	re options
(Do	ctor to document in space provided.	Continue ii
Med	lical Record if necessary.)	

E. Risks of not having this procedure

(Doctor to document in space provided. Continue in Medical Record if necessary.)

F Anaesthetic	
	_
	•

This procedure may require an anaesthetic. (Doctor to document type of anaesthetic discussed)

Queensland Government
(Government

Fractured Forearm

	(Amx identification labe	i nere)		
URN:				
Family name:				
Given name(s):				
Address:				
Date of birth:		Sex: M	F	

Facility:

G. Patient consent

I acknowledge that the doctor has explained;

- my medical condition and the proposed procedure, including additional treatment if the doctor finds something unexpected. I understand the risks, including the risks that are specific to
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- other relevant procedure/treatment options and their associated risks.
- my prognosis and the risks of not having the procedure.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- the procedure may include a blood transfusion.
- tissues and blood may be removed and could be used for diagnosis or management of my condition, stored and disposed of sensitively by the hospital.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor or my Acute Resuscitation Plan.
- a doctor other than the Consultant may conduct the procedure. I understand this could be a doctor undergoing further training.

I have been given the following Patient Information Sheet/s:

About Your Anaesthetic OR
Anaesthetic: Nerve Blocked

- Fractured Forearm
- I was able to ask questions and raise concerns with the doctor about my condition, the proposed procedure and its risks, and my treatment options. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time, including after I have signed this form but, preferably following a discussion with my doctor.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.

On the basis of the above statements,

I request to have the procedure

Name of Pa	tient:
Signature:	
Date:	
Patients	s who lack capacity to provide consent
	must be obtained from a substitute decision nthe order below.
Does the (AHD)?	patient have an Advance Health Directive
☐ Yes ►	Location of the original or certified copy of the AHD:
□ No ►	Name of Substitute Decision Maker/s: Signature:
	Relationship to patient:
	Date: PH No:
	Source of decision making authority (tick one):
	Tribunal-appointed Guardian
	Attorney/s for health matters under Enduring Power of Attorney or AHD
	☐ Statutory Health Attorney
	If none of these, the Adult Guardian has provided consent. Ph 1300 QLD OAG (753 624)
L	

H. Doctor/delegate Statement

I have explained to the patient all the above points under the Patient Consent section (G) and I am of the opinion that the patient/substitute decision-maker has understood the information.

Name of Doctor/delegate:	
Designation:	
Signature:	
Date:	

I. Interpreter's statement

I	have	given	а	sight	trans	lation	ir
---	------	-------	---	-------	-------	--------	----

(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

guardian/substitute decision-maker by the doctor.
Name of
Interpreter:
Signature:
Date:



Consent Information - Patient Copy Fractured Forearm

What do I need to know about this procedure?

This procedure is the fixation of the fractured forearm with the placement of plate and screws or nails.

2. My anaesthetic

This procedure will require an anaesthetic.

See About Your Anaesthetic OR Anaesthetic: **Nerve Block** for information about the anaesthetic and the risks involved. If you have any concerns, discuss these with your doctor.

If you have not been given an information sheet, please ask for one.

3. What are the risks of this specific procedure?

There are risks and complications with this procedure. They include but are not limited to the following.

- · Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

Specific risks:

- · Numbness associated with the use of tourniquet with nerve and muscle damage at the site where the tourniquet was placed. This may be temporary or permanent.
- Skin death under the tourniquet, which may require further dressings and / or surgery and skin grafting.
- Damage to nerves and/or vessels resulting in numbness and/or weakness. This may temporary or permanent. Another operation may be required to correct the situation.
- Stiffness in the joints may occur above and below the break in the arm.
- Build up of pressure in the muscle compartment, which can require re-operation.
- Inability to close the edges of the wound together. This may require skin grafts.

- The bones do not knit together properly. This may require further surgery and/or bone graft.
- The bones heal with a bend in the bones and/or alignment.
- The implants may need to be removed in future months or years. This will require further surgery.
- Abnormal pain response to surgery with worsening of pain and disability.
- The surgical cut may cause changes to the sensation and colour of the limb.
- In some people, healing of the wound may be abnormal and the wound can be thickened and red and the scar may be painful.

Notes	to talk to	my docto	r about: