

National Immunisation Program Queensland Schedule

February 2018

CHILDHOOD SCHEDULE [^] Caution – be aware of brand name similarity. Give all vaccinations on time as scheduled.

Age	Antigen	Vaccine Brand	Route	Comments
Birth	Hepatitis B	H-B-Vax II® (paediatric) [^] OR Engerix®B (paediatric)	IM IM	Within 7 days of birth – ideally within the first 24 hours
6 weeks (42 days)	DTPa-hep B-IPV-Hib Pneumococcal Rotavirus	Infanrix hexa® [^] Prevenar 13® (13vPCV) Rotarix®	IM IM Oral	Observe Rotarix® administration age limits: Dose 1: 6-14 weeks of age Dose 2: 10-24 weeks of age
4 months	DTPa-hep B-IPV-Hib Pneumococcal Rotavirus	Infanrix hexa® [^] Prevenar 13® (13vPCV) Rotarix®	IM IM Oral	
6 months	DTPa-hep B-IPV-Hib Pneumococcal	Infanrix hexa® [^] Prevenar 13® (13vPCV)	IM IM	
12 months	MMR Hib-Meningococcal C	M-M-R II® OR Priorix® Menitorix®	S/C S/C or IM IM	Additional vaccines are recommended for children with medical risk factors and/or children born prematurely (see below)
Additional vaccines for Aboriginal and Torres Strait Islander children	Hepatitis A	Vaqta® Paediatric	IM	
18 months	MMRV DTPa	Priorix Tetra® [^] OR ProQuad® Infanrix® [^] OR Tripacel®	S/C or IM S/C IM	
Additional vaccines for Aboriginal and Torres Strait Islander children	Pneumococcal Hepatitis A	Prevenar 13® (13vPCV) Vaqta® Paediatric	IM IM	
4 years	DTPa-IPV	Infanrix IPV® [^] OR Quadracel®	IM	Additional vaccines are recommended for children with medical risk factors (see below)

ADOLESCENT / ADULT SCHEDULE

Year 7 Students	HPV	Gardasil®9	IM	<ul style="list-style-type: none"> • 2 dose course given at 0 and at least 6 months between dose 1 and dose 2 • Immunocompromised individuals require 3 doses of Gardasil®9 at 0, 2 and 6 months (refer to the immunocompromised conditions listed in Additional notes provided with this schedule)
	dTpa	Boostrix®	IM	Single dose
Year 10 Students and 15 to 19 year olds	Meningococcal ACWY	Menactra®	IM	The Meningococcal ACWY Program will continue through 2018 for Year 10 students and 15 to 19 year olds through immunisation providers
Pregnant women (From 28 weeks gestation)	dTpa	Adacel®	IM	Single dose recommended each pregnancy. Administer between 28-32 weeks gestation for optimal protection
Aboriginal & Torres Strait Islander adults 15 - 49 years	Pneumococcal	Pneumovax 23® (23vPPV)	IM	For further information please refer to the online version of The Australian Immunisation Handbook at www.health.gov.au/internet/immunise/publishing.nsf/Content/Handbook10-home~handbook10part4~handbook10-4-13
Aboriginal & Torres Strait Islander adults 50 years and over				
65 years and over	Pneumococcal	Pneumovax 23® (23vPPV)	IM	
70 years of age	Varicella Zoster (shingles)	Zostavax®	S/C	Single dose
71-79 years of age	Varicella Zoster (shingles)	Zostavax®	S/C	A single catch-up dose funded until 2021

Additional funded vaccines for medically at risk individuals

12 months	Hepatitis B: Children born at <32 weeks gestation and/or <2000g birth weight are recommended to be given a booster dose of hepatitis B vaccine at 12 months of age Pneumococcal (13vPCV): Children with medical risk factors including premature infants born at <28 weeks gestation are recommended to be given a 4th dose of 13vPCV at 12 months of age
4 years	Pneumococcal (23vPPV): Children with medical risk factors including premature infants born at <28 weeks gestation are recommended to be given a booster dose of 23vPPV vaccine at 4-5 years of age
Any age group	Pneumococcal (23vPPV): Adults with medical risk factors may require additional 23vPPV vaccine

For further information refer to the online version of the *Australian Immunisation Handbook* www.immunise.health.gov.au

Refer to the current National Influenza Immunisation Program resources

For more information contact your local public health unit

Cairns	4226 5555	Rockhampton	4920 6989	Brisbane North	3624 1111	Brisbane South	3176 4000
Townsville	4433 6900	Wide Bay	4303 7500	Sunshine Coast	5409 6600	West Moreton	3818 4700
Mt Isa	4744 7186	Mackay	4885 5800	Gold Coast	5667 3200	Darling Downs	4699 8240