Clinical Excellence Queensland

GLYCEROL (GLYCERINE) SUPPOSITORY

Indication		 Promote meconium passage¹ Short-term management of constipation² 					
RECTAL	Presentation	Suppository 700 mg					
	Dosage ³	Single dose once					
		Current gest age (weeks)*	Dose				
		Less than 32+0	Small chip (less than one quarter)				
		32+0 to 39+6	Quarter suppository				
		40+0 to 43+6	One half to one full suppository				
		44+0 or more	Full suppository				
2		*Current gestational age is t	the same as postmenstrual age (PMA)				
	Preparation	 Remove suppository from packaging and hold firmly with sterile metal forceps Using a sterile blade, cut lengthwise to obtain prescribed dose size Round off sharp/abrasive cut edges by smoothing with gloved finger 					
	Administration	 Moisten suppository with lubricant⁴ Insert fully into rectum³ 					
	Special considerations	 Preterm infant Dosage recommendations (limited evidence) based on consensus and clinical judgement for gestational age and weight⁴ No increased risk of rectal perforation or NEC^{1,5} 					
	Monitoring	For passage of meconium or bowel motions					
	Compatibility	Nil known					
	Incompatibility	Nil known					
	Interactions	nteractions • Nil significant					
	Stability	 Store at room temperature below 30 °C Discard unused portion 					
	Side effects	Digestive: abdominal discomfort ³ , local irritation ⁶					
	Actions	 Osmotic laxative with lubricating properties⁶ May act as a stimulant by its local irritant effects⁷ Onset of action 15–60 minutes⁸ 					
	Abbreviations	Current gestational age is the same as postmenstrual age (PMA) NEC: necrotising enterocolitis					
	Keywords neonatal medicine, neonatal monograph, suppository, constipation, laxative, aperient, peristalsis, NEC, necrotising enterocolitis, meconium						

The Queensland Clinical Guideline *Neonatal Medicines* is integral to and should be read in conjunction with this monograph. Refer to the disclaimer. Destroy all printed copies of this monograph after use.



References

- 1. Burchard PR, Lay R, Ruffolo LI, Ramazani SN, Walton JM, Livingston MH. Glycerin suppositories and enemas in premature infants: a meta-analysis. Pediatrics 2022;149(4).
- 2. British National Formulary for Children (BNFC) online. Glycerol. [Internet]: Royal Pharmaceutical Society; November 2021 [cited 2023 March 30]. Available from: https://www.medicinescomplete.com.
- 3. MIMS Online. Petrus Glycerol Suppositories BP. [Internet]: MIMS Australia; March 2023 [cited 2023 March 30]. Available from: https://www.mimsonline.com.au.
- 4. Jannin V, Lemagnen G, Gueroult P, Larrouture D, Tuleu C. Rectal route in the 21st century to treat children. Advanced Drug Delivery Reviews 2014;73:34-49.
- 5. Gross M, Hummler H, Haase B, Quante M, Wiechers C, Poets CF. Interventions for promoting meconium passage in very preterm infants-a survey of current practice at tertiary neonatal centers in Germany. Children (Basel) 2022;9(8).
- 6. Anabrees J, Shah VS, AlOsaimi A, AlFaleh K. Glycerin laxatives for prevention or treatment of feeding intolerance in very low birth weight infants. Cochrane Database of Systematic Reviews. [Internet]. 2015, [cited 2023 March 30]. Issue 9. Art No.: CD010464. DOI:10.1002/14651858.CD010464.pub2.
- 7. Australian Medicines Handbook. Glycerol. [Internet]. Adelaide: Australian Medicines Handbook Pty Ltd; January 2023 [cited 2023 April 12]. Available from: https://amhonline.amh.net.au.
- 8. Australian Medicines Handbook Children's Dosing Companion. Glycerol. [Internet]. Adelaide: Australian Medicines Handbook Pty Ltd; January 2023 [cited 2023 March 30]. Available from: https://www.amhonline.amh.net.au.

Document history

ID number	Effective	Review	Summary of updates
NMedQ23.108-V1-R28	22/09/2023	22/09/2028	Endorsed by Queensland Neonatal Services Group (QNSAG)

QR code

