

Specific Catch-up Recommendations for Gardasil 9[®] Vaccine

Doses of Gardasil 9 [®] recommended if adolescent currently aged ≤14 years (i.e. < 15 years)	
No. of Gardasil (HPV) doses given previously	Doses of Gardasil 9 [®] (HPV9) recommended
No previous doses	Two doses needed, give at least 6 months apart Give at 0, 6-12 months
1 previous dose	One dose needed, give at least 6 months after previous HPV dose. <i>No further doses required as 2 doses will complete the schedule</i>
2 previous doses <i>Note: doses 1 & 2 should be given at least 6 months apart. If dose 2 was given < 6 months and ≥5 months no further doses are required.</i>	None, if dose 1 & 2 were given at least 6 months apart <ul style="list-style-type: none"> • <u>However, if dose 2 was given less than 5 months after dose 1</u> <ul style="list-style-type: none"> – Give dose 3 at least 12 weeks after dose 2 and at least 5 months after dose 1, whichever is later
3 previous doses <i>Note: Revaccination with a course of Gardasil 9 is not recommended or funded for those who have previously completed a funded course of Gardasil (HPV) vaccine</i>	None, if minimum spacing between all 3 doses are met, at least: <ul style="list-style-type: none"> • 4 weeks between dose 1 & 2 • 12 weeks between dose 2 & 3 and • 5 months between dose 1 & 3

Immunocompromised Adolescents

All adolescents who are immunocompromised due to the specific conditions listed below require a three-dose course of Gardasil 9 – given at 0, 2, 6 months.

Primary or secondary immune-deficiencies: B lymphocyte antibody & T lymphocyte complete or partial deficiencies, HIV infection, Malignancy, Organ transplantation, significant immunosuppressive therapy.

Doses of Gardasil 9 [®] recommended if adolescent currently aged ≥ 15 years	
No. of Gardasil (HPV) doses given previously	Doses of Gardasil 9 (HPV9) recommended
No previous doses	Three doses needed. Give at 0, 2, 6 months
1 previous dose given <15 years	One dose needed, give at least 6 months after previous HPV dose. <i>No further doses required as 2 doses will complete the schedule</i>
1 previous dose given ≥15 years	Two doses needed, <ul style="list-style-type: none"> • give dose 2 at least two months after dose 1 • give dose 3 at least four months after dose 2
2 previous doses given <15 years <i>Note: doses 1 & 2 should be given at least 6 months apart. If dose 2 was given < 6 months and ≥5 months no further doses are required.</i>	None, if dose 1 & 2 were given at least 6 months apart <ul style="list-style-type: none"> • <u>However, if dose 2 was given less than 5 months after dose 1</u> <ul style="list-style-type: none"> – Give dose 3 at least 12 weeks after dose 2 and at least 5 months after dose 1, whichever is later
2 previous doses given ≥15 years	One dose needed, give at least four months after dose 2
3 previous doses given at any age <i>Note: Revaccination with a course of Gardasil 9 is not recommended or funded for those who have previously completed a funded course of Gardasil (HPV) vaccine</i>	None, if minimum spacing between all 3 doses are met, at least: <ul style="list-style-type: none"> • 4 weeks between dose 1 & 2 • 12 weeks between dose 2 & 3 and • 5 months between dose 1 & 3

This content is accurate at the date of approval. Contact the Sunshine Coast Public Health Unit Tel 1300 017 190 for further information.