



**Queensland
 Government**

Lumbar Puncture for Intrathecal Chemotherapy Consent

Facility:

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

A. Does the patient have capacity to provide consent?

Complete for ADULT patient only

- Yes → **GO TO section B**
 No → **COMPLETE section A**

You must adhere to the Advance Health Directive (AHD), or if there is no AHD, the consent obtained from a substitute decision-maker in the following order: Category 1. Tribunal-appointed guardian; 2. Enduring Power of Attorney; or 3. Statutory Health Attorney.

Name of substitute decision-maker:

Category of substitute decision-maker:

Complete for CHILD/YOUNG PERSON patient only

- Yes Although the patient is a child/young person, the patient may be capable of giving informed consent and having sufficient maturity, understanding and intelligence to enable them to fully understand the nature, consequences and risks of the proposed procedure and the consequences of non-treatment – ‘Gillick competence’ (*Gillick v West Norfolk and Wisbech Area Health Authority* [1986] AC 112)
 → **GO TO section B**
- No Parent/legal guardian/other person* with parental rights and responsibilities to provide consent and complete this form
 → **COMPLETE section A**

*Formal arrangements, such as parenting/custody orders, adoption, or other formally recognised carer/guardianship arrangements. Refer to the Queensland Health ‘Guide to Informed Decision-making in Health Care’ and local policy and procedures. Complete the source of decision-making authority as applicable below.

If applicable, source of decision-making authority (*tick one*):

- Court order → Court order verified
 Legal guardian → Documentation verified
 Other person → Documentation verified

Name of parent/legal guardian/other person:

Relationship to child/young person:

B. Is an interpreter required?

- Yes No

If yes, the interpreter has:

- provided a sight translation of the informed consent form in person
 translated the informed consent form over the telephone

It is acknowledged that a verbal translation is usually a summary of the text on the form, rather than word-by-word translation.

Name of interpreter:

Interpreter code:

Language:

C. Patient OR substitute decision-maker OR parent/legal guardian/other person confirms the following procedure(s)

I confirm that the referring doctor/clinician has explained that I have been referred for the following procedure:

Lumbar puncture for intrathecal chemotherapy: Yes No

Name of referring doctor/clinician:

D. Risks specific to the patient in having a lumbar puncture for intrathecal chemotherapy

(Doctor/clinician to document additional risks not included in the patient information sheet):

E. Risks specific to the patient in not having a lumbar puncture for intrathecal chemotherapy


(Doctor/clinician to document specific risks in not having a lumbar puncture for intrathecal chemotherapy):

F. Alternative procedure options

(Doctor/clinician to document alternative procedure not included in the patient information sheet):

DO NOT WRITE IN THIS BINDING MARGIN

v1.00
 Clinical content review: 2023
 Clinical check: 07/2023
 Published: 07/2023



SW9611

LUMBAR PUNCTURE FOR INTRATHECAL CHEMOTHERAPY CONSENT



Lumbar Puncture for Intrathecal Chemotherapy Consent

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

G. Information for the doctor/clinician

The information in this consent form is not intended to be a substitute for direct communication between the doctor/clinician and the patient OR substitute decision-maker OR parent/legal guardian/other person.

I have explained to the patient OR substitute decision-maker OR parent/legal guardian/other person the contents of this form and am of the opinion that the information has been understood.

Name of doctor/clinician:

Designation:

Signature:

Date:

H. Patient OR substitute decision-maker OR parent/legal guardian/other person consent

I acknowledge that the doctor/clinician has explained:

- the 'Lumbar Puncture for Intrathecal Chemotherapy' patient information sheet
- the medical condition and proposed treatment, including the possibility of additional treatment
- the specific risks and benefits of the procedure
- the prognosis, and risks of not having the procedure
- alternative procedure options
- that there is no guarantee the procedure will improve the medical condition
- that tissues/blood may be removed and used for diagnosis/management of the condition
- that if a life-threatening event occurs during the procedure:
 - an adult patient will be treated based on documented discussions (e.g. AHD or ARP [Acute Resuscitation Plan])
 - a child/young person's health care will be provided in accordance with good clinical practice and in the best interests of the patient
- that a doctor/clinician other than the consultant/specialist may assist with/conduct the clinically appropriate procedure; this may include a doctor/clinician undergoing further training under supervision
- that if the doctor/clinician wishes to record video, audio or images during the procedure where the recording is not required as part of the treatment (e.g. for training or research purposes), I will be asked to sign a separate consent form. If I choose not to consent, it will not adversely affect my access, outcome or rights to medical treatment in any way.

I was able to ask questions and raise concerns with the doctor/clinician.

I understand I have the right to change my mind regarding consent at any time, including after signing this form (*this should be in consultation with the doctor/clinician*).

I/substitute decision-maker/parent/legal guardian/other person have received the following consent and patient information sheet(s):

- 'Lumbar Puncture for Intrathecal Chemotherapy'
- 'About Your Anaesthetic' (*Adult patient only*)
- 'About Your Child's Anaesthetic' (*Child/young person patient only*)

On the basis of the above statements,

1) I/substitute decision-maker/parent/legal guardian/other person consent to having a lumbar puncture for intrathecal chemotherapy.

Name of patient/substitute decision-maker/parent/legal guardian/other person:

Signature:

Date:

If the patient is a child/young person:

- I am not aware of any legal or other reason that prevents me from providing unrestricted consent for this child/young person for this procedure (*not applicable if the child/young person is Gillick competent and signs this form*).

2) Student examination/procedure for professional training purposes:

For the purpose of undertaking training, a clinical student(s) may observe medical examination(s) or procedure(s) and may also, subject to patient OR substitute decision-maker OR parent/legal guardian/other person consent, assist with/conduct an examination or procedure on a patient while the patient is under anaesthetic.

I/substitute decision-maker/parent/legal guardian/other person consent to a clinical student(s) undergoing training to:

- observe examination(s)/procedure(s) Yes No
- assist with examination(s)/procedure(s) Yes No
- conduct examination(s)/procedure(s) Yes No

Lumbar Puncture for Intrathecal Chemotherapy

Adult and Child/Young Person | Informed consent: patient information

A copy of this patient information sheet should be given to the patient or substitute decision-maker or parent/legal guardian/other person of a child or young person to read carefully and allow time to ask any questions about the procedure. The consent form and patient information sheet should be included in the patient's medical record.

In this information sheet, the word 'you' means the patient unless a substitute decision-maker, parent, legal guardian or other person is providing consent on behalf of the patient, in which case the word 'you' means the substitute decision-maker, parent, legal guardian or other person when used in the context of the person providing consent to the procedure.




1. What is a lumbar puncture for intrathecal chemotherapy and how will it help me?

A lumbar puncture for intrathecal chemotherapy injection is a procedure that uses imaging to guide a needle into position, so chemotherapy drugs can be injected into your spinal fluid to treat tumour cells in the fluid, or sometimes when there is a high risk of tumour spread to the fluid.

Sometimes, a small amount of Cerebrospinal Fluid (CSF) will be collected and sent to pathology for testing.

*Image: Lumbar puncture. Illustration Copyright © 2019 Nucleus Medical Media, All rights reserved.
www.nucleusmedicalmedia.com*



Preparing for the procedure

The Medical Imaging department will give you instructions on how to prepare for the procedure. It is important to follow the instructions that are given to you. Your procedure might be delayed if you don't follow all the preparation steps.

Medical imaging staff will notify you beforehand if you are required to stop taking any blood-thinning medicine. List or bring all your prescribed medicines, those medicines you buy over the counter, herbal remedies and supplements to show the doctor/clinician what you are taking.

It is important that you lie still for the procedure. If unable to lie still, adults may be given sedation and a child or young person may require a general anaesthetic. These would need to be organised prior to the date of your procedure.

If you received sedation or a general anaesthetic and are being discharged on the same day, you cannot drive and you must have someone available to escort you home.

Do not drink alcohol, smoke, vape or take recreational drugs for at least 24 hours before the procedure as these may alter the effects of the sedation anaesthetic. If you have a drug or medication dependence, please tell your doctor/clinician.

Please tell the doctor/clinician if you are breastfeeding or pregnant, or suspect that you may be pregnant.

On the day of your procedure

- Nothing to eat or drink ('nil by mouth'): you will be told when to have your last meal and drink. Do NOT eat (including lollies), drink or chew gum after this time otherwise your procedure may be delayed or cancelled. This is to make sure your stomach is empty so that if you vomit, there will be nothing to go into your lungs.
- If you are taking medicines, most should be continued before a procedure and taken at the usual time, even on the day of the procedure, with a sip of water. There are some important exceptions:
 - your doctor/clinician will provide specific instructions about your medicines
 - take to the hospital all your prescribed medicines, those medicines you buy over the counter, herbal remedies and supplements to show your doctor/ nurse what you are taking.
- If you feel unwell, telephone the medical imaging department for advice
- Tell your doctor/clinician if you have:
 - health problems (e.g. diabetes, high blood pressure, infectious diseases, serious illnesses), including if undergoing regular treatment
 - a drug or medicine dependence
 - had previous problems and/or known family problems with anaesthesia
 - been taking prescribed and/or over the counter medicines, herbal remedies and supplements; this may include and is not limited to blood thinning medicines, the contraceptive pill, antidepressants and/or medicine for treating diabetes (e.g. insulin)
 - allergies/intolerances of any type and their side effects
- You may/will be required to change into a hospital gown and remove some of your jewellery.

Sedation

Sedation is the use of medicines that help make you feel relaxed and drowsy for your procedure. You may remember some or little about what has happened. You may still be aware of your surroundings and should be able to follow simple instructions, such as holding your breath when instructed by the doctor/clinician.

If you are booked for an anaesthetic or sedation, please read the information sheet *About Your Anaesthetic (for adults)* or *About Your Child's Anaesthetic (for child/young person)*. If you do not have one of these information sheets, please ask for one.

For a parent/legal guardian/other person of a patient having a lumbar puncture for intrathecal chemotherapy

To prepare the patient for this procedure and to ease their concerns, tell them what they can expect to happen during the procedure. This information sheet will assist you with this.

We welcome your help and support in preparing the patient for the procedure and in explaining why it's so important to lie still.

At the discretion of the procedure staff:

- a parent/adult (unless pregnant) may be invited into the procedure room to support the patient
- if the patient is having a general anaesthetic you may be able to see them off to sleep. Once they are asleep you will be asked to leave the procedure room and wait in the waiting area.

Other children are not allowed into the procedure room, and they must be supervised at all times by another parent/adult.

During the procedure

You will be positioned on the examination table lying on your front or side. Supporting straps, foam pads and light weights may also be used to help support you.

An intravenous (I.V.) cannula is a small plastic tube that will be inserted into a vein, usually in your hand or arm, if sedation or a general anaesthetic is required.

The skin on your lower back will be cleaned and a sterile drape will be applied to cover you.

A local anaesthetic is injected into the skin and soft tissues of your lower back. Local anaesthetic will numb the area and is used to prevent and relieve pain but will not put you to sleep. A spinal needle is guided into the space between two of your back bones until it reaches the spinal canal.

While the needle is inserted try not to move or cough suddenly. X-ray images are taken to help guide the needle into the spinal canal.

When the needle is in the spinal canal, an oncology doctor will inject the chemotherapy drugs through the needle and into your spinal fluid.

The needle is then removed from your back and a dressing is applied.

You will need to lie flat for a period of time after the procedure. This is to reduce the risk of developing a headache.

You will need to take care when first standing as temporary leg weakness can occur.

If you had an I.V. cannula and it is no longer required, it will be removed.



2. What are the risks?

In recommending the procedure, the doctor/clinician believes that the benefits to you from having the procedure exceed the risks involved. There are risks and possible complications associated with the procedure which can occur with all patients – these are set out below. There may also be additional risks and possible complications specific to your condition and circumstances which the doctor/clinician will discuss with you. If you have any further concerns, please ensure that you raise them with the doctor/clinician prior to giving consent to the procedure.

Common risks and complications

- headache, this may require medicine and bed rest
- minor pain, bruising and/or infection at the I.V. cannula site or the injection site on your back. This may require treatment

- bleeding/bruising is more common if you have been taking blood thinning medicines, such as warfarin, aspirin, clopidogrel (Plavix, Iscover, Coplavix), prasugrel (Effient), dipyridamole (Persantin or Asasantin), ticagrelor (Brilinta), apixaban (Eliquis), dabigatran (Pradaxa), rivaroxaban (Xarelto) or complementary/alternative medicines, such as fish oil and turmeric
- low blood pressure causing dizziness. Bed rest can help relieve this
- failure of local anaesthetic which may require a further injection of anaesthetic or a different method of anaesthesia may be used
- nerve damage, which is usually temporary and should get better over time.

Uncommon risks and complications

- the procedure may not be possible due to medical or technical reasons
- a severe headache which may resolve with a few days of bed rest. Sometimes other procedures are required to be done to try to relieve the headache
- infection requiring antibiotics and further treatment
- damage to surrounding structures such as blood vessels, organs and muscles, requiring further treatment
- an allergy to injected drugs, requiring further treatment
- vomiting may occur and require treatment.

Rare risks and complications

- injury to the spinal cord. This may require surgery
- permanent nerve damage with possible paralysis
- meningitis requiring antibiotics and other treatment
- seizures and/or cardiac arrest due to local anaesthetic toxicity.
- death because of this procedure is very rare.

If general anaesthetic or sedation is given, risks include:

- faintness or dizziness, especially when you start to move
- fall in blood pressure
- nausea and vomiting
- weakness

- heart and lung problems, such as heart attack or pneumonia
- stroke resulting in brain damage.

Risks of radiation

The risks of radiation exposure from this procedure need to be compared to the risks of your condition not being treated. Exposure to radiation may cause a slight increase in the risk of cancer to you over your lifetime. However, the potential risk is small compared to the expected benefit of this procedure¹.

What are the risks of not having a lumbar puncture for intrathecal chemotherapy?

There may be adverse consequences for your health if you choose not to have the proposed procedure. Please discuss these with the referring doctor/clinician.

If you choose not to have the procedure, you will not be required to sign a consent form.

If you have signed a consent form, you have the right to change your mind at any time prior to the procedure. Please contact the doctor/clinician to discuss.



3. Are there alternatives?

Making the decision to have a procedure requires you to understand the options available. Please discuss any alternative procedure options with your doctor/clinician before signing the consent form.



4. What should I expect after the procedure?

Your healthcare team will talk to you about what to expect after your procedure and upon discharge from hospital and what level of activity is suitable.

Go to your nearest Emergency department or GP if you become unwell or have:

- light sensitivity
- agitation
- amnesia
- seizures
- pain unrelieved by simple pain relievers

- continuous bleeding or swelling at the puncture site
- redness or inflammation at the puncture site
- fever
- other warning signs the doctor/clinician may have asked you to be aware of.

If you had sedation, this will affect your judgement for about 24 hours. For your own safety:

- Do NOT drive any type of car, bike or other vehicle.
- Do NOT operate machinery including cooking equipment.
- Do NOT make important decisions or sign a legal document.
- Do NOT drink alcohol, smoke, vape or take recreational drugs. They may react with the anaesthetic medications.



5. Who will be performing the procedure?

Doctors, radiographers, nuclear medicine technologists, sonographers, nurses, and medical imaging assistants make up the medical imaging team. All or some of these professionals may be involved in your procedure.

An oncologist and other oncology clinicians will participate in this procedure.

A doctor/clinician other than the consultant/specialist may assist with/conduct the clinically appropriate procedure. This could include a doctor/clinician undergoing further training, however all trainees are supervised according to relevant professional guidelines.

If you have any concerns about which doctor/clinician will be performing the procedure, please discuss this with the doctor/clinician.

For the purpose of undertaking professional training in this teaching hospital, a clinical student(s) may observe medical examination(s) or procedure(s) and may also, subject to your consent, assist with/conduct an examination or procedure on a patient while the patient is under anaesthetic.

You are under no obligation to consent to an examination(s) or a procedure(s) being undertaken by a clinical student(s) for training purposes. If you choose not to consent, it will not adversely affect your access, outcome or rights to medical treatment in any way.

For more information on student care, please visit www.health.qld.gov.au/consent/students.



6. Where can I find support or more information?

Hospital care: before, during and after is available on the Queensland Health website www.qld.gov.au/health/services/hospital-care/before-after where you can read about your healthcare rights.

You can also see a list of blood thinning medications at www.health.qld.gov.au/consent/bloodthinner.

Further information about informed consent can be found on the Informed Consent website www.health.qld.gov.au/consent. Additional statewide consent forms and patient information sheets are also available here.

Staff are available to support patients' cultural and spiritual needs. If you would like cultural or spiritual support, please discuss this with your doctor/clinician.

Queensland Health recognises that Aboriginal and Torres Strait Islander patients will experience the best clinical care when their culture is included during shared decision-making.



7. Questions

Please ask the doctor/clinician if you do not understand any aspect of this patient information sheet or if you have any questions about your proposed procedure.

If you have further questions prior to your appointment, please contact the Medical Imaging department via the main switchboard of the facility where your procedure is booked.



8. Contact us

In an emergency, call Triple Zero (000).

If it is not an emergency, but you have concerns, contact 13 HEALTH (13 43 25 84), 24 hours a day, 7 days a week.

References:

1. Australian Radiation Protection and Nuclear Safety Agency (ARPANSA). Ionising radiation in our everyday environment, 2021. Available from www.arpansa.gov.au