



Queensland Government

Coiling of Intra-Cranial Aneurysm

Facility:

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

A. Interpreter / cultural needs

- An Interpreter Service is required? Yes No
If Yes, is a qualified Interpreter present? Yes No
A Cultural Support Person is required? Yes No
If Yes, is a Cultural Support Person present? Yes No

B. Procedure

The following will be performed (*Doctor/doctor delegate to document – include site and/or side where relevant to the procedure*)

.....

.....

An Intra-Cranial Aneurysm is an abnormality and weakness in the wall of a blood vessel in the brain.

Coiling (or embolisation) is where tiny coils of soft metal thread are used to block off the blood flow in the aneurysm.

Coils are inserted into the aneurysm the blood inside the aneurysm clots and prevents any more blood entering it. This reduces the chance of the aneurysm bursting or bleeding. The coils stay in for life.

Depending on the shape of your aneurysm a stent may be required. A stent is a small metal mesh tube that is placed in the artery to help hold the coils inside your aneurysm and prevent them from falling out. Aneurysms that have wide necks usually need a stent. If used, the stent also stays in for life.

This procedure will require the injection of a local anaesthetic and a general anaesthetic.

C. Risks of the procedure

In recommending the Coiling of Intra-Cranial Aneurysm the doctor believes the benefits to you from having this procedure exceeds the risks involved.

The risks and complications with this procedure can include but are not limited to the following.

Common risks and complications include:

- Minor pain, bruising and/or infection from the IV cannula. This may require treatment with antibiotics.
- Pain or discomfort at the puncture site. This may require medication.
- Bleeding or bruising may occur. This is usually stopped by applying pressure and/or ice to the puncture site. This is more common if you take Aspirin, Warfarin, Clopidogrel (Plavix and Iscover) or Dipyridamole (Persantin and Asasantin).
- Failure of local anaesthetic which may require a further injection of anaesthetic or a different method of anaesthesia may be used.

- Nerve damage, is usually temporary, and should get better over a period of time. Permanent nerve damage is rare.

Less common risks and complications include:

- Infection, requiring antibiotics and further treatment.
- Rupture of the aneurysm causing bleeding in the brain. This can lead to stroke or stroke like complications which can cause weakness in the face, arms and legs. This could be temporary or permanent.
- Stroke caused by constricted blood vessels, blood clots or ruptured artery. This can cause weakness in the face, arms and legs. This could be temporary or permanent.
- Damage to surrounding structures such as blood vessels, organs and muscles, requiring further treatment.
- A blood clot or excessive bleeding from the puncture site. This may require other treatment and/or corrective surgery.
- An allergy to injected drugs, requiring further treatment.
- Over time, the compressed coils may cause the aneurysm to reform. This may require further coiling or surgical procedures.
- The procedure may not be possible due to medical and/or technical reasons.

Rare risks and complications include:

- An increased lifetime cancer risk due to the exposure to x-rays.
- Skin burns or damage from exposure to x-rays.
- Incomplete coiling of your aneurysm. This may require further coiling procedures.
- Seizures and/or cardiac arrest due to local anaesthetic toxicity.
- Death as a result of this procedure is possible.

D. Risks of Iodinated Contrast for patients with renal impairment

Specific Risks of Iodinated Contrast to patient's identified as having *Renal Impairment*.

- Giving the Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).



Coiling of Intra-Cranial Aneurysm

Facility:

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex: M F I

E. Patient consent

I acknowledge that the doctor/doctor delegate has explained the proposed procedure.

I understand;

- the risks and complications, including the risks that are specific to me.
- the sedation/ anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor/doctor delegate or my Acute Resuscitation Plan.
- a doctor/doctor delegate undergoing further training may conduct this procedure.

I have been given the following Patient Information Sheet/s:

- Coiling of Intra-Cranial Aneurysm
- Iodinated Contrast
- About Your Anaesthetic

- I was able to ask questions and raise concerns with the doctor/doctor delegate about the proposed procedure and its risks. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time including after I have signed this form but, preferably following a discussion with my doctor/doctor delegate.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.
- I understand that Queensland Health may release my relevant de-identified information obtained from this and related procedures for education and training of health professionals.

On the basis of the above statements,

I request to have the procedure

Name of Patient:.....

Signature:.....

Date:.....

Patients who lack capacity to provide consent

Consent must be obtained from a substitute decision maker/s in the order below.

Does the patient have an Advance Health Directive (AHD)?

Yes ▶ Location of the original or certified copy of the AHD:.....

No ▶ Name of Substitute Decision Maker/s:.....
Signature:.....
Relationship to patient:.....
Date:..... PH No:.....

Source of decision making authority (tick one):

- Tribunal-appointed Guardian
- Attorney/s for health matters under Enduring Power of Attorney or AHD
- Statutory Health Attorney
- If none of these, the Adult Guardian has provided consent. Ph 1300 QLD OAG (753 624)

F. Doctor/delegate Statement

I have explained to the patient all the above points under:

- the Patient Consent section (E)
- Iodinated Contrast - Patients with Renal Impairment Section (D) (for renal impaired patients only)

and I am of the opinion that the patient/substitute decision-maker has understood the information.

Name of Doctor/delegate:.....

Designation:.....

Signature:..... Date.....

G. Interpreter's statement

I have given a sight translation in

.....
(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

Name of Interpreter:.....

Signature:..... Date.....


DO NOT WRITE IN THIS BINDING MARGIN

Consent Information - Patient Copy

Coiling of Intra-Cranial Aneurysm

1. What is a Coiling of an Intra-Cranial Aneurysm?

An Intra-Cranial Aneurysm is an abnormality and weakness in the wall of a blood vessel in the brain.




Aneurysm, Blood-Flow - Herston Multi Media Unit, RBWH, 2009

Coiling (or embolisation) is where tiny coils of soft metal thread are used to block off the blood flow in the aneurysm.

Filling the aneurysm with coils stops the blood flowing into it, reducing the chance of the aneurysm bursting and bleeding. When coils are inserted into the aneurysm the blood inside the aneurysm clots and prevents any more blood entering it. The coils stay in for life.


Coiling is used instead of open surgery to treat aneurysms.



Coiling - Herston Multi Media Unit, RBWH, 2009

Stents

Depending on the shape of your aneurysm a stent may be required. A stent is a small metal mesh tube that is placed in the artery to help hold the coils inside your aneurysm and prevent them from falling out. Aneurysms that have wide necks usually need a stent. If used, the stent also stays in for life.



Coiled Aneurysm

The Coiling is done by placing a needle and a thin plastic tube (catheter) into the artery in your groin. Iodinated 'Contrast' (once called x-ray dye) is used during the procedure to map your arteries so the doctor can locate and treat your aneurysm. For more information on Iodinated Contrast and the risks involved in its use, please read the **Iodinated Contrast Patient Information Sheet**. (If you do not have this information sheet please ask for one).

2. Will there be any discomfort, is any anaesthetic needed?

This procedure will require both a local and general anaesthetic. For more information on general anaesthetics and the risks involved, please read the **About your Anaesthetic Patient Information sheet**. (If you do not have this information sheet please ask for one).

3. Preparation for the procedure

The medical imaging department will give you instructions on how to prepare for your procedure.

- You may be given blood thinning medications to take for a few days before your appointment.
- Please tell the staff if you are or suspect you might be pregnant or are breastfeeding.

4. During the procedure

A fine needle (IV cannula) will be put into a vein in your arm.

You will be given a general anaesthetic.

The Radiologist (x-ray doctor) will inject local anaesthetic into the skin. A needle and catheter are inserted into the artery in your groin; sometimes your arm may be used. Once the catheter is in place the needle is removed.

The catheter is guided through the main blood vessels in your body until it reaches the blood vessels that go to your brain.

X-ray pictures are taken while the Contrast is injected into your arteries.

If required, a stent will be placed in the artery across the opening of the aneurysm.

Once the catheter is in place the coils are gently pushed into the aneurysm. The coils wind and bend conforming to the shape of the aneurysm slowly filling it up. Several coils may be needed to completely fill your aneurysm.

Contrast is injected after every coil is placed to assess your aneurysm.

Once the aneurysm is filled with coils the catheter is removed leaving the coils in place.

Firm pressure will be put over the area where the catheter went into your skin (puncture site), sometimes a special plug is used. This allows the artery to seal so you will not bleed.

1. What is Iodinated Contrast?

The medical imaging procedure your doctor has asked you to have uses Iodinated 'Contrast' (once called X-ray dye). Contrast is a colourless liquid which includes iodine. Contrast is injected into your blood stream to allow your organs to be seen more clearly on x-rays. Contrast is not a dye. It does not stain the inside of your body. Your doctor needs to use Contrast to be able to get all the information needed to assist with your diagnosis.

This information sheet must be read together with the information sheet of the procedure you are booked for (if you do not have this information sheet please ask for one).

2. During the procedure

When the Contrast is injected you may feel:

- A very warm or 'flushed' feeling over your body, this may also make you think you have passed urine. You will *not* pass urine – it is only a feeling.
- A 'metallic' taste or smell may also happen. This usually lasts less than a minute.

3. After the procedure

It is recommended that you drink 2 to 4 glasses of water after your procedure to help flush the Contrast from your body.

Contrast does not affect your ability to carry out normal activities; you should be able to continue with your day as normal.

4. Precautions

Contrast is not suitable for some people; you will be asked a series of questions before it is given to you. Your answers allow staff to identify any risk factors that you may have.

- Please tell the staff if you are or suspect you might be pregnant or are breastfeeding.

Kidney function

- Contrast is removed from your blood by your kidneys through your urine. It is easily removed from the body of people who have normal kidney function.
- Giving Contrast to people with weakened kidneys (renal impairment), can cause further kidney damage, which may in turn cause the kidneys to stop working properly (acute renal failure).
- You may be asked to have a simple blood test to find out the level of their kidney function.

Diabetic Drug interactions - Metformin

(Other Drug names: Avandamet, Diabex, Diaformin, Formet, Glucohexal, Glucomet, Glucophage, Glucovance, Metforbell)

If kidneys suffer damage from the Contrast then the kidneys may not be able to remove Metformin from the body. It is safer to briefly stop taking Metformin when having Contrast. Staff will inform you when to stop and when it is safe to take Metformin again.

Contact your GP to monitor your diabetes if you are told to stop your Metformin.

5. What are the risks of Iodinated Contrast?

The risks and complications with this injection can include but are not limited to the following.

Common risks and complications include:

- No known common risks.

Less common risks and complications include:

- Injected Contrast may leak outside of the blood vessel, under the skin and into the tissue. This may require treatment. In very rare cases, further surgery could be required if the skin breaks down.
- Acute Renal Failure occurs when one or both of your kidneys suddenly stop working. Failure can last for days or weeks. It may take the kidneys a long time to regain their previous level of function and you may require dialysis to filter your blood during this time. There is a risk your kidneys could be permanently damaged. To reduce this risk the smallest possible dose of Contrast will be given.
- The injection may not be possible due to medical and/or technical reasons.

Rare risks and complications include:

- Allergic reactions occur within the first hour with most happening in the first 5 minutes. Late reactions have been known to occur up to a week after the injection.

Note: Allergy to topical iodine and/or seafood does not imply an allergy to Iodinated Contrast.

The reactions vary from:

Mild – hives, sweating, sneezing, coughing, nausea.

Moderate – wide spread hives, headaches, facial swelling, vomiting, shortness of breath.

Severe – Severe reactions are rare but include: life-threatening heart palpitations, very low blood pressure, throat swelling, fits and/or cardiac arrest.

- Death as a result of Iodinated Contrast is *very* rare.

6. What are the safety issues when you leave the hospital?

Go to your nearest Emergency Department or GP if you become unwell.