

In-patient Falls Assessment and Management Plan

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Adult

Sex: M F I

Facility:

- Complete assessment within eight (8) hours of admission
- Reassess at a minimum of weekly, when there is a change in condition, medication, after a fall and on discharge
- Care plans never replace clinical judgement. Care outlined must be altered if it is not clinically appropriate for the individual patient
- Every person documenting on the form must supply a sample of their initials in the signature log (page2)

Falls Risk Assessment

Identify risk factors Tick (✓) Yes or No <i>(if Yes to any, patient is 'at risk' of a fall)</i>	If YES to any →	Initiate actions Tick when actioned (if indicated)
--	--------------------	--

Risk Factors	Date	Time	Initial		Date	Time	Initial
Screen: The patient has had a fall in the last 6 months	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Refer patient to physiotherapist for gait and balance assessment		
The patient is observed to be unsteady	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient requires supervision or assistance with transfer	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Conduct pre-activity screening prior to off bed transfer		
The patient is visually impaired	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient has new onset incontinence	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Initiate ward urinalysis • Notify MO and facilitate tests as ordered (e.g. MSU)		
The patient has existing incontinence, frequency or requires assisted toileting	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient reports postural symptoms	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Measure lying and standing BP		
The patient has a recent history of syncope	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient is on one of the following medications: (antihypertensive, antidepressant, sedative, antipsychotic, benzodiazepine)	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Refer to MO / Pharmacist for medication review / simplification		
The patient is on more than 4 medications	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient has a minimal trauma fracture and / or history of osteoporosis	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Facilitate tests ordered by MO (e.g. TFT, calcium, vitamin D assay, PTH, sEPP) • Refer to Dietitian (as appropriate)		
The patient has new onset or increased confusion / delirium	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				
The patient is usually confused	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	▶	• Conduct or refer for cognitive assessment (if appropriate)		
	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N	<input type="checkbox"/> Y <input type="checkbox"/> N				

Following assessment, proceed to management plan for ALL patients (page 2)

DO NOT WRITE IN THIS BINDING MARGIN





**Queensland
Government**

In-patient Falls Assessment and Management Plan

(Affix identification label here)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Adult

Sex: M F I

- **Complete within eight (8) hours of admission**
- Review management plan at a minimum daily and document as per local policy
- Initial when strategies are implemented
- **V** indicates a variance from clinical care and must be documented in the clinical notes

Falls Prevention Management Plan

All care givers who initial are to sign signature log

⌘ Key ◆ Allied Health ■ Medical ▲ Nursing ⊕ Pharmacy

Category	⌘		Date		
			Time		
Communication	▲	In partnership with patient and / or carer discuss falls risk factors and develop falls prevention plan			
	◆	Provide written and verbal falls prevention information (e.g. <i>Stay On Your Feet® BE SAFE brochure</i>)			
	■	using the Australian Commission on Safety and Quality in Health Care (ACSQHC) teach back method			
	⊕	Communicate patients 'at risk' status at bedside handover			
		Instruct patient to call for assistance when getting out of bed / mobilising (if required)			
		Falls risk patients on anti-coagulant / antiplatelet medication, request MO / Pharmacy medication review			
Environment / Equipment <i>(used according to local policy and procedures)</i>	▲	Orientate patient to surroundings, routine and location of bathroom and toilet			
		Ensure clutter free and safe environment (e.g. night time lighting)			
		Ensure the chair and bed height / position are suitable for the patient's needs (i.e. the patient's feet need to be on the ground with the knees slightly below the hip)			
		Apply brakes to bed, wheelchair and commode correctly			
		Ensure use of bed rails are appropriate for patient's needs and if used, height is appropriate			
		Keep buzzer in reach; educate patient on buzzer usage			
	Keep patient's routine belongings within reach				
	Keep patient's current mobility aid assessed as safe to use within reach (where applicable), remove all other mobility aids from patient's sight and reach				
	Review patient footwear and / or foot problems				
Other Care <i>(specify)</i>	▲				
	◆				
	■				
	⊕				
Observations	▲	Ensure frequent rounding and surveillance			
		Consider supervision during toileting / showering / mobilisation			
		Ensure suitable toileting protocols are in place			
Specific Patient Centred Goal <i>(e.g. prefers to wear closed in shoes when transferring / mobilising)</i>	▲				
	◆				
	⊕				
	■				
Discharge Planning / Education	▲	Provide information on falls risk factors and prevention strategies (e.g. <i>Stay On Your Feet® Checklist</i>)			
	◆	Refer to OT for ADL and home assessment			
		Complete nursing discharge summary and facilitate referrals			

DO NOT WRITE IN THIS BINDING MARGIN

Signature Log (Every person documenting on the form [page 1 and 2] must supply a sample of their initials)

Initial	Print name	Designation	Signature	Initial	Print name	Designation	Signature