























# Queensland Health Immunisation Program

## NIP Funded Pneumococcal Vaccines (From 1 July 2026)

### Product Selection Guide

This product selection guide is to be used in conjunction with the Immunisation Schedule Queensland and the Australian Immunisation Handbook.

Australian Immunisation Handbook - Pneumococcal chapter 	Immunisation Schedule Queensland 	Adult pneumococcal schedule program advice 	<b>Prevenar 20*</b>  20-valent pneumococcal conjugate vaccine	<b>Capvaxive†</b>  21-valent pneumococcal conjugate vaccine
<b>Infants ≤12 months</b> 	<b>All infants**</b>	 Dose 1 at age 2 months (can be from 6 weeks) Dose 2 at age 4 months Dose 3 at age 12 months	 Not registered for use	
	<b>Aboriginal and Torres Strait Islander infants AND/OR infants with a medical risk condition**</b>	 Additional dose at age 6 months	 Not registered for use	
<b>Children and adolescents &lt;18 years</b> 	<b>Children and adolescents aged &gt;12 months to &lt;18 years with a newly diagnosed medical risk condition**</b>	 Single dose at diagnosis	 Not registered for use	
	<b>Aboriginal and Torres Strait Islander people AND/OR people with a medical risk condition at 4 years of age OR 5 years after first Pneumovax 23 dose**</b>	 Single dose ONLY required if child has not previously received a dose of Prevenar 20	 Not registered for use	
<b>Adults ≥18 years</b> 	<b>Adults ≥18 years with a newly diagnosed medical risk condition*</b>	 Not funded for use	 Single dose at diagnosis	
	<b>Aboriginal and Torres Strait Islander people ≥25 years</b>	 Not funded for use	 Single dose	
	<b>All adults ≥65 years</b>	 Not funded for use	 Single dose	

Only 1 dose of Capvaxive recommended in a person's lifetime. Ensure Capvaxive is administered at least 12 months after any previous pneumococcal vaccine.

\*Prevenar 20 is the only pneumococcal vaccine funded for individuals under the age of 18 years.

PneumococcalProductSelectionGuide\_V3\_22062026

†Capvaxive is the only pneumococcal vaccine funded for individuals aged 18 years and over.

\*Refer to the Australian Immunisation Handbook for more information.