



QUEENSLAND OLDER ADULT IMMUNISATION VISUAL TOOL




QUICK REFERENCE GUIDE TO FUNDED VACCINES AND ELIGIBILITY CRITERIA

ADULTS AGED 65 YEARS AND OVER AND ABORIGINAL AND TORRES STRAIT ISLANDER ADULTS AGED 50 YEARS AND OVER



18 YEARS AND OVER	Shingles					
	Shingrix  2 doses, 1-2 months apart in people who are immunocompromised or shortly expected to be immunocompromised.	Medically at risk 	50 YEARS AND OVER	Shingrix  2 doses, 2-6 months apart.	Aboriginal and Torres Strait Islander 	65 YEARS AND OVER
	Shingrix 2 doses, 2-6 months apart.					


Pneumococcal						
18 YEARS AND OVER	Capvaxie  1 dose at least 12 months after any previous pneumococcal vaccine.	Medically at risk	25 YEARS AND OVER	Capvaxie  1 dose at least 12 months after any previous pneumococcal vaccine.	Aboriginal and Torres Strait Islander	65 YEARS AND OVER
						
	Capvaxie 1 dose at least 12 months after any previous pneumococcal vaccine.					
						

20 YEARS AND OVER	Diphtheria, Tetanus and Pertussis (Whooping Cough)		Hepatitis B			
	Boostrix  1 booster dose if >10 years (>5 years in high-risk areas) since last diphtheria vaccine. If a person has not completed a primary course, vaccines are funded regardless of age.	Aboriginal and Torres Strait Islander 	HB-Vax II Adult or Engerix B Adult Funded for at risk people aged 20 years and over who have not previously received a valid hepatitis B vaccine course Use Engerix-B only for people living with HIV. Use HB-Vax II Adult for other eligible people.	Aboriginal and Torres Strait Islander	Medically at risk	At-risk populations
						

COVID-19							
50 TO 74 YEARS	Comirnaty  1 dose annually.	Aboriginal and Torres Strait Islander	Comirnaty 1 dose annually.		75 YEARS AND OVER	Comirnaty 2 doses per year (approximately 6 months apart).	
							

60 YEARS AND OVER	Respiratory syncytial virus (RSV)			Influenza		
	Arexvy  1 dose.	Aboriginal and Torres Strait Islander 	75 YEARS AND OVER	Arexvy 1 dose.		
				LESS THAN 65 YEARS	Vaxigrip OR Flucelvax 1 dose annually. 	65 YEARS AND OVER
						Fluad 1 dose annually. 

ALL AGES	Meningococcal ACWY		Meningococcal B		Haemophilus influenzae type b (Hib)	
	Nimenrix  2 doses at least 8 weeks apart for people with specified medical risk conditions. Ongoing booster doses may be required.	Medically at risk 	Bexsero  2 doses at least 8 weeks apart for people with specified medical risk conditions.	Medically at risk 	Act-HIB  1 dose for people with asplenia or hyposplenia who were not vaccinated in childhood.	Medically at risk Act-HIB Haemophilus type b Conjugate Vaccine (conjugated to tetanus protein) Haemophilus type b polysaccharide 10 microgram (conjugated to tetanus protein 18-30 microgram) Powder and diluent for solution + 5 mL of diluent in pre-filled syringe with 2 separate needles Intramuscular or deep subcutaneous injection AUST R 130161 

Japanese Encephalitis Virus				Mpox	
9 MONTHS AND OVER	Imojev  1 dose.	Funded for eligible at-risk people in specified Queensland locations, including people with increased mosquito exposure. Check current location and eligibility criteria.	ALL AGES	JEspect  2 doses at least 28 days apart. Only funded for immunocompromised people, during pregnancy and infants aged 2 to <9 months.	Funded for eligible people at risk of exposure. Check current eligibility criteria.
					Jynneos  2 doses at least 28 days apart. 

Key:  Aboriginal and Torres Strait Islander people  Medically at risk  At-risk populations



This visual tool is accurate at the time of publication and supports, but does not replace, appropriate clinical assessment. Always use in conjunction with the Australian Immunisation Handbook, the Immunisation Schedule Queensland and current Queensland Health program advice.

V1 - 28092026 © State of Queensland (Queensland Health) 2026. Contact: immunisation@health.qld.gov.au

DELIVERING
FOR QUEENSLAND



Queensland
Government