











QUEENSLAND ADOLESCENT AND ADULT IMMUNISATION VISUAL TOOL

QUICK REFERENCE GUIDE TO FUNDED VACCINES AND ELIGIBILITY CRITERIA

ADOLESCENTS AND ADULTS AGED LESS THAN 65 YEARS AND ABORIGINAL AND TORRES STRAIT ISLANDER ADULTS LESS THAN 50 YEARS							
YEAR 7 (OR AGE EQUIVALENT)	Human Papillomavirus (HPV)			Diphtheria, Tetanus and Pertussis (Whooping Cough)			
	Gardasil 9 1 dose. A 3-dose schedule is recommended for persons with certain immunocompromising conditions.			Boostrix OR Adacel 1 dose.			
YEAR 10 (OR AGE EQUIVALENT)	Meningococcal ACWY			Meningococcal B			
	MenQuadfi 1 dose. Funded for Year 10 students (or age equivalent) AND adolescents aged 15-19 years inclusive.			Bexsero 2 doses, at least 8 weeks apart. Funded for Year 10 students (or age equivalent) AND adolescents aged 15-19 years inclusive.			
LESS THAN 18 YEARS	Pneumococcal						
	Prevenar 20 1 dose at diagnosis of specified medical risk condition.		18 YEARS AND OVER	Capvaxime 1 dose at least 12 months after any previous pneumococcal vaccine.		25 YEARS AND OVER	Capvaxime 1 dose at least 12 months after any previous pneumococcal vaccine.
18 YEARS AND OVER	Shingles			COVID-19			
	Shingrix 2 doses, 1-2 months apart in people who are immunocompromised or shortly expected to be immunocompromised.		18 TO 74 YEARS	Comirnaty 1 dose annually in people with severe immunocompromise.			
20 YEARS AND OVER	Diphtheria, Tetanus and Pertussis (Whooping Cough)			Hepatitis B			
	Boostrix 1 booster dose if >10 years (>5 years in high-risk areas) since last diphtheria vaccine. If a person has not completed a primary course, vaccines are funded regardless of age.		Aboriginal and Torres Strait Islander	HB-Vax II or Engerix B Adult Funded for people aged 20 years and over who are at increased risk and have not previously completed a valid hepatitis B vaccine course. Engerix-B available for people living with HIV. HB-Vax II Adult for other eligible people.			
ALL PEOPLE BORN DURING OR SINCE 1966	Measles, mumps, rubella (MMR)			LESS THAN 65 YEARS	Influenza		
	Priorix OR M-M-RII 2 doses at least 4 weeks apart. Funded for people without documented evidence of 2 doses of measles-containing vaccine.		Vaxigrip OR Flucelvax 1 dose annually.				
ALL AGES	Meningococcal ACWY		Meningococcal B		Haemophilus influenzae type b (Hib)		
	Nimenrix 2 doses at least 8 weeks apart for people with specified medical risk conditions. Ongoing booster doses may be required.		Bexsero 2 doses at least 8 weeks apart for people with specified medical risk conditions. Ongoing booster doses may be required.		Act-HIB 1 dose for people with asplenia or hyposplenia who were not vaccinated in childhood.		
9 MONTHS AND OVER	Japanese Encephalitis Virus				Mpox		
	Imojev 1 dose.		JEspect 2 doses at least 28 days apart. Only funded for immunocompromised people, during pregnancy and infants aged 2 to <9 months.		Jynneos 2 doses at least 28 days apart.		
DURING PREGNANCY							
DURING PREGNANCY	Influenza		20-32 WEEKS PREGNANT	Pertussis (Whooping cough)		FROM 28 WEEKS PREGNANT	Respiratory Syncytial Virus (RSV)
	Vaxigrip OR Flucelvax 1 dose at any time in each pregnancy.			Boostrix OR Adacel 1 dose in each pregnancy between 20-32 weeks, but may be given up until birth.			