

Preterm prelabour rupture of membranes (PPROM)

Clinical Guideline Presentation v1.0



45 minutes

Towards CPD Hours

References:

Queensland Clinical Guideline: Preterm prelabour rupture of membranes is the primary reference for this package.

Recommended citation:

Queensland Clinical Guidelines. Preterm prelabour rupture of membranes clinical guideline education presentation E18.48-1-V1-R23. Queensland Health 2018.

Disclaimer:

This presentation is an implementation tool and should be used in conjunction with the published guideline. This information does not supersede or replace the guideline. Consult the guideline for further information and references.

Feedback and contact details:

M: GPO Box 48 Brisbane QLD 4001 | **E:** guidelines@health.qld.gov.au | **URL:** www.health.qld.gov.au/qcg

Funding:

Queensland Clinical Guidelines is supported by the Queensland Health, Healthcare Improvement Unit.

Copyright:

© State of Queensland (Queensland Health) 2018



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International licence. In essence, you are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute Queensland Clinical Guidelines, Queensland Health and abide by the licence terms. You may not alter or adapt the work in any way. To view a copy of this licence, visit <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For further information, contact Queensland Clinical Guidelines, RBWH Post Office, Herston Qld 4029, email Guidelines@health.qld.gov.au, phone (07) 3131 6777. For permissions beyond the scope of this licence, contact: Intellectual Property Officer, Queensland Health, GPO Box 48, Brisbane Qld 4001, email ip_officer@health.qld.gov.au, phone (07) 3234 1479.

Images are property of State of Queensland (Queensland Health) unless otherwise cited.

Objectives

- Identify management options following PPROM
- Identify risks and benefits of management options



Preterm PROM

What is PPROM?

PPROM refers to women who:

- Have ruptured membranes
- Are less than 37+0 weeks gestation
- Are not in labour

Preterm Prelabour Rupture Of Membranes

- Occurs in around 3% of pregnancies
- 50% go into labour within one week
- 75% go into labour within two weeks
- 25-50% have an infection at presentation

Diagnosis

Avoid digital vaginal examinations as may increase risk of infection

How is the diagnosis of PPRM made?

- History:
 - Sudden gush or continued leakage of fluid from vagina
- Sterile speculum:
 - Pooling of amniotic fluid or leakage from cervical os with coughing
- Test vaginal secretions:
 - Immunoassay (e.g. Amnisure®)
 - pH stick (e.g. Nitrazine)

Conduct a clinical assessment

- Review history
- Maternal vital signs
- General physical exam
- Abdominal palpation
- Fetal heart rate/CTG

During speculum exam

- Exclude cord prolapse
- Visualise cervical length and dilatation



Initial assessment

Isla is 33 weeks pregnant with her first baby. She thinks her 'waters have broken'. After clinical assessment you confirm PPRM. She is otherwise well.

What additional investigations will you recommend?

- Routine antenatal bloods not already collected
- Full blood count +/- baseline CRP
- Dipstick urinalysis MSU for MC&S
- Swabs for GBS (may be self-collected)
- With sterile speculum collect high vaginal swab
- Ultrasound for fetal wellbeing and to confirm presentation

Aim of care

Maximise benefits of increasing fetal maturity in-utero while minimising potential risks

CRP

Conflicting evidence about usefulness in PPRM—may be a useful adjunct to exclude infection as has low sensitivity (less than 20%) and high specificity (greater than 90%) for chorioamnionitis

Antibiotics

Isla asks whether she should have antibiotics.

Should you recommend antibiotics?

Yes. Antibiotics are known to prolong latency and reduce maternal and fetal infection following PPRM

Avoid

Amoxicillin/clavulanic acid as associated with increased risk of necrotising enterocolitis in baby



Recommend either

- Erythromycin 250 mg oral 6 hourly for 10 days OR
- Amoxicillin/ampicillin 2 g IV every 6 hours for 48 hours, followed by amoxicillin 250 mg oral every 8 hours for a total of 7 days (IV and oral), PLUS erythromycin 250 mg oral every 6 hours for 7 days

Inpatient care

You recommend Isla be admitted to hospital for 72 hours for assessment and monitoring.

In addition to monitoring for infection, what care will you give Isla?

- Counsel about prematurity
- Involve neonatologist/paediatrician in care planning
- Offer psychological support
- Involve other members of the multidisciplinary team relevant to circumstances (e.g. social worker, maternal fetal medicine specialist, Aboriginal and Torres Strait Islander health worker)

Self care advice

- Advise about the risk of infection
- Personal hygiene—change pad four hourly (or more frequently)
- Wipe front to back after toileting
- Showering in preference to baths
- Avoiding tampon use, vaginal creams/medications, vaginal intercourse, immersion in water (e.g. swimming, bathing, spa)
- Attending all review appointments

Risk of infection

Isla asks about the risk of infection and asks how she will know if she gets one?

What advice do you give Isla?

- The risk of infection is increased after PPRM
- Report any concerns
- Self-monitor for signs of infection and report same
- Follow self-care advice
- Attend all appointments (if discharged)



Signs of infection

- Feeling unwell or flu-like symptoms
- Maternal temperature greater than 37.5 °C
- Change in vaginal discharge (odour, volume, amount, colour)
- Uterine tenderness
- Fetal tachycardia
- Change in fetal movements

Outcomes

Isla asks about the risks and benefits of planned birth if she were to have her baby now.

What can you tell Isla about planned birth at 33 weeks after PPROM?

No difference in:

- Chorioamnionitis
- Caesarean section
- Neonatal infection

Decreased

- Endometritis



If PPROM, after 34 weeks with any planned birth versus expectant care:

Decreased

- Chorioamnionitis
- APH

Increased

- Respiratory distress

Birth

Isla asks when is the best time to have her baby.

What can you tell Isla about this?

- Optimal time is unknown
- Individual circumstances will determine timing
- Dependent on ongoing health and wellbeing of mother and baby
- Generally, less than 34 weeks, favours waiting



If preterm birth:

- Antenatal corticosteroids if under 35 weeks
- Magnesium sulfate for neuroprotection if under 30 weeks
- In-utero transfer if limited service capacity
- Recommend intrapartum GBS prophylaxis